

Recommendations for competing sexual-asexually typified generic names in *Sordariomycetes* (except *Diaporthales*, *Hypocreales*, and *Magnaporthales*)

Martina Réblová¹, Andrew N. Miller², Amy Y. Rossman^{3*}, Keith A. Seifert⁴, Pedro W. Crous⁵, David L. Hawksworth^{6,7,8}, Mohamed A. Abdel-Wahab⁹, Paul F. Cannon⁸, Dinushani A. Daranagama¹⁰, Z. Wilhelm De Beer¹¹, Shi-Ke Huang¹⁰, Kevin D. Hyde¹⁰, Ruvvishika Jayawardena¹⁰, Walter Jaklitsch^{12,13}, E. B. Gareth Jones¹⁴, Yu-Ming Ju¹⁵, Caroline Judith¹⁶, Sajeewa S. N. Maharachchikumbura¹⁷, Ka-Lai Pang¹⁸, Liliane E. Petrini¹⁹, Huzefa A. Raja²⁰, Andrea I. Romero²¹, Carol Shearer², Indunil C. Senanayake¹⁰, Hermann Voglmayr¹³, Bevan S. Weir²², and Nalin N. Wijayawarden¹⁰

¹Department of Taxonomy, Institute of Botany of the Academy of Sciences of the Czech Republic, Průhonice 252 43, Czech Republic

²Illinois Natural History Survey, University of Illinois, Champaign, Illinois 61820, USA

³Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331, USA; *corresponding author e-mail: amydianer@yahoo.com

⁴Ottawa Research and Development Centre, Biodiversity (Mycology and Microbiology), Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, Ontario K1A 0C6 Canada

⁵CBS-KNAW Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands

⁶Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense, Plaza de Ramón y Cajal s/n, Madrid 28040, Spain

⁷Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

⁸Comparative Plant and Fungal Biology, Royal Botanic Gardens, Kew, Surrey, TW9 3DS, UK

⁹Department of Botany and Microbiology, Faculty of Science, Sohag University, Sohag 82524, Egypt

¹⁰Center of Excellence in Fungal Research, School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

¹¹Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria 0002, South Africa

¹²Institute of Forest Entomology, Forest Pathology and Forest Protection, Department of Forest and Soil Sciences, BOKU-University of Natural Resources and Life Sciences, Vienna, Austria

¹³Division of Systematic and Evolutionary Botany, Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria

¹⁴Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

¹⁵Institute of Plant and Microbial Biology, Academia Sinica, Nankang, Taipei 115 29, Taiwan

¹⁶Department of Mycology, Institute of Ecology, Evolution and Diversity, Goethe-University, Max-von-Laue-Str. 13, 60438 Frankfurt am Main, Germany

¹⁷Department of Crop Sciences, College of Agricultural and Marine Sciences, Sultan Qaboos University, PO Box 8, 123 Al Khoud, Oman

¹⁸Institute of Marine Biology and Centre of Excellence for the Oceans, National Taiwan Ocean University, 2 Pei-Ning Road, Keelung 20224, Taiwan (ROC)

¹⁹Via al Perato 15c, CH-6932 Breganzona, Switzerland

²⁰Department of Chemistry and Biochemistry, 457 Sullivan Science Building, University of North Carolina, Greensboro, NC 27402-6170, USA

²¹Instituto de Micología y Botánica, UBA-CONICET, Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Pabellón II, Piso 4°, Lab 6, Av. Int. Güiraldes 2620. Ciudad Universitaria, C1428EHA, Buenos Aires, Argentina

²²Manaaki Whenua Landcare Research, Private Bag 92170, Auckland, New Zealand

Abstract: With the advance to one scientific name for each fungal species, the generic names in the class *Sordariomycetes* typified by sexual and asexual morphs are evaluated based on their type species to determine if they compete with each other for use or protection. Recommendations are made for which of the competing generic names should be used based on criteria such as priority, number of potential names changes, and frequency of use. Some recommendations for well-known genera include *Arthrinium* over *Apiospora*, *Colletotrichum* over *Glomerella*, *Menispora* over *Zignoëlla*, *Microdochium* over *Monographella*, *Nigrospora* over *Khuskia*, and *Plectosphaerella* over *Plectosporium*. All competing generic names are listed in a table of recommended names along with the required action. If priority is not accorded to sexually typified generic names after 2017, only four names would require formal protection: *Chaetosphaerella* over *Oedemium*, *Diatrype* over *Libertella*, *Microdochium* over *Monographella*, and *Phaeoacremonium* over *Romellia* and *Togninia*. Concerning species in the recommended genera, one replacement name (*Xylaria benjaminii* nom. nov.) is introduced, and the following new combinations are made: *Arthrinium sinense*, *Chloridium caesium*, *C. chloroconium*, *C. gonytrichii*, *Corollospora marina*, *C. parvula*, *C. ramulosa*, *Juncigena fruticosae*, *Melanospora simplex*, *Seimatosporium massarina*, *Sporoschisma daemnoropis*, *S. taitense*, *Torpedospora mangrovei*, *Xylaria penicillioopsis*, and *X. termiticola* combs. nov.

Key words:

Ascomycota
nomenclature
pleomorphic fungi
protected lists
taxonomy

Article info: Submitted: 26 April 2016; Accepted: 24 May 2016; Published: 8 June 2016.

© 2016 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

INTRODUCTION:

The class *Sordariomycetes* is composed of three subclasses and about 21 orders including many genera with species that express themselves in both their sexual and asexual morphs. Based on the obsolete version of Article 59 of the outdated *International Code of Botanical Nomenclature* (McNeill et al. 2006), these morphs had previously been described in different genera resulting in more than one scientific name for a single fungal species. With the change to the *International Code of Nomenclature for algae, fungi, and plants* (ICN; McNeill et al. 2012), two or more names for different morphs of the same species are no longer allowed. Although determining which name to use generally follows the principle of priority of publication at the family, generic and species level, exceptions to this principle are allowed, especially in the case of economically important and widely used taxa.

In this paper generic names that appear to compete for use are reviewed to determine if their respective type species are congeneric using various resources including the USDA SMML Fungal Databases (<http://nt.ars-grin.gov/fungaldatabases/>) and Wijayawardene et al. (2012). If so, then a number of factors are considered in deciding which generic name to recommend for use. These factors include the number of species in each competing genus, which correlates with the number of name changes that would be required, and how widely used are species in each genus as determined by reports and peer-reviewed publications. If these factors are about equal, then the generic name that has priority by date is recommended for use. For each set of competing generic names, these factors are discussed based on the literature. Finally, a draft of these recommendations was circulated widely amongst the community of mycologists interested in each major group of fungi to arrive at the proposals made here.

A synopsis of data concerning each genus is provided in Table 1 listing the generic names recommended for use and competing names, the type species for each genus and the current name for the type, and action required especially if the principle of priority will not be followed, i.e. the generic name must be protected. At present the ICN requires that names typified by sexual morphs should have priority unless protected, although it is proposed that this requirement be deleted in 2017 (Hawksworth 2015). Eventually all generic names proposed for protection will be evaluated by the Nomenclature Committee for Fungi (NCF), and formally accepted or not at the Nomenclature Section meeting of the next International Botanical Congress in 2017.

For several groups of fungi having pleomorphic genera, this process has been completed, i.e. *Dothideomycetes* (Rossman et al. 2015b), *Erysiphales* (Braun 2013), *Eurotiales* (Samson et al. 2014, Visagie et al. 2014), *Leotiomyces* (Johnston et al. 2014), yeast fungi (Daniel et al. 2014), and some orders of *Sordariomycetes* including *Diaporthales* (Rossman et al. 2015a), *Hypocreales* (Rossman et al. 2013, Quandt et al. 2014), *Magnaporthales* (Zhang et al. 2016), and *Microascales* and *Ophiostomatales* (de Beer et al. 2013). This paper addresses the remaining orders of the class *Sordariomycetes* listed alphabetically by order as outlined in Maharachchikumbura et al. (2015): *Amphisphaeriales*,

Calosphaeriales, *Chaetosphaeriales*, *Coniochaetales*, *Conioscyphales*, *Cordanales*, *Coronophorales*, *Glomerellales*, *Halosphaeriales*, *Melanosporales*, *Microascales*, *Phyllachorales*, *Pleurotheciales*, *Savoryellales*, *Sordariales*, *Togniniales*, *Torpedosporales*, and *Xylariales*.

RECOMMENDATIONS FOR GENERIC NAMES

(A) = a name typified by an asexual morph, and (S) = a name typified by a sexual morph.

Amphisphaeriales

Use *Dyrithiopsis* L. Cai et al. 2003 (S) rather than *Monochaetiopsis* L. Cai et al. 2003 (A)

The monotypic genera *Dyrithiopsis*, typified by *D. lakefuxianensis*, and *Monochaetiopsis*, typified by *M. lakefuxiansis*, were described as the sexual and asexual morph of the same species at the same time (Jeewon et al. 2003), thus they are synonyms and have equal priority. Because *Dyrithiopsis* has been cited more widely in the literature, we recommend *Dyrithiopsis* for use.

Use *Hyalotiopsis* Punith. 1970 (A) rather than *Ellurema* Nag Raj & W.B. Kendr. 1985 (S)

When Punithalingam (1970) described the genus *Hyalotiopsis*, typified by *H. subramaniani*, he noted that the sexual morph was *Massaria indica*, the basionym of *Ellurema indica*, type of the monotypic *Ellurema*, thus *Hyalotiopsis* and *Ellurema* are synonyms. This history was recounted by Nag Raj & Kendrick (1986) when they described *Ellurema*. A second species of *Hyalotiopsis* has been redisposed in *Parahyalotiopsis* as *P. borassi* (syn. *Hyalotiopsis borassi*) (Nag Raj 1976), thus both genera include only one species. Given their equal use, we follow priority and recommend *Hyalotiopsis* for use.

Use *Pestalotiopsis* Steyaert 1949 (A) rather than *Pestalosphaeria* M.E. Barr 1975 (S)

The well-known genus *Pestalotiopsis*, typified by *P. guepinii*, was established for many species of *Pestalotia* determined to be distinct from the type of *Pestalotia*, *P. pezizoides*. Barr (1975) described the genus *Pestalosphaeria*, typified by *P. concentrica*, for the sexual morph of *Pestalotiopsis guepinii* var. *macrotricha*, both species on *Rhododendron* in southeastern North America. Thus, it appears that *Pestalosphaeria* is a synonym of *Pestalotiopsis*. *Pestalotiopsis* has been widely used and currently includes more than 200 names with two recent accounts of the genus (Maharachchikumbura et al. 2012, 2014). On the other hand, *Pestalosphaeria* includes 14 names, none of which are commonly used. Given the ubiquitous and widely reported species of *Pestalotiopsis* as well as the number of names in that genus and its priority, we recommend use of *Pestalotiopsis*.

Use *Seimatosporium* Corda 1833 (A) rather than *Discostroma* Clem. 1909 (S)

The generic names *Discostroma* and *Seimatosporium* have been applied to sexual and asexual morphs of the same species based on morphological studies (Brockmann

1976, Nag Raj 1993). The type species of *Discostroma*, *D. rehmsii*, is a younger synonym of *D. massarina*, and has an asexual morph referred to as *Seimatosporium ribis-alpini* while the type of *Seimatosporium*, *S. rosae*, has a sexual morph described as *Discostroma rosae* (Shoemaker 1964, Brockmann 1976, Nag Raj 1993). Although neither of the type species of *Discostroma* or *Seimatosporium* were included, Tanaka *et al.* (2011) used three species of *Discostroma* and sixteen species of *Seimatosporium* in their LSU tree to show that representatives of these genera form a monophyletic genus that should be regarded as *Seimatosporium*. Norphanphoun *et al.* (2015) added four more isolates of *Seimatosporium* including one for the type species, *S. rosae*, and also concluded that the genus was monophyletic. Host-specificity should be considered in studying the taxonomy of this genus. Given that *Seimatosporium* is the oldest name, has the greater number of species, and is more commonly used, this generic name is recommended for use.

New combination:

***Seimatosporium massarina* (Sacc.) Jaklitsch & Voglmayr, comb. nov.**

MycoBank MB817259

Basionym: *Metasphaeria massarina* Sacc., *Atti Soc. Veneto-Trent. Sci. Nat., Padova*, Sér. 6 2: 22 (1884).

Synonyms: *Discostroma massarina* (Sacc.) Arx, *Gen. Fungi Sporul. Cult.*, 2nd edn. (Vaduz): 131 (1974).

Hendersonia ribis-alpini Fautrey, *Revue Mycol.*, Toulouse 14: 171 (1892).

Seimatosporium ribis-alpini (Fautrey) Shoemaker & E. Müll., *Canad. J. Bot.* 42: 403 (1964).

Curreya rehmsii Schnabl, *Ber. Bayer. Bot. Ges., Beih.* 2: 66 (1892)

Discostroma rehmsii (Schnabl) Clem., *Gen. Fungi*: 173 (1909).

Use *Seiridium* Nees 1816 (A) rather than *Blogiascospora* Shoemaker *et al.* 1966 (S)

According to Shoemaker *et al.* (1966) *Seiridium marginatum*, the type species of *Seiridium*, is the asexual morph of *Blogiascospora marginata*, type of the monotypic genus *Blogiascospora*, thus *Seiridium* and *Blogiascospora* are synonyms as confirmed by Jaklitsch *et al.* (2016). *Seiridium* is the older generic name, has the greater number of species, and is the more widely used, therefore *Seiridium* is recommended for use.

One species of *Seiridium*, *S. cupressi*, cause of the widespread disease known as cypress canker, has previously been referred to as *Lepteutypa cupressi*. The type of *Lepteutypa*, *L. fuckelii*, is now placed in a distinct genus in the *Amphisphaeriaceae* (Jaklitsch *et al.* 2016), while the species previously regarded as *L. cupressi* belongs in *Seiridium* as *S. cupressi* in *Sporocadaceae*.

Calosphaeriales

Use *Calosphaeria* Tul. & C. Tul. 1863 (S) rather than *Calosphaeriophora* Réblová *et al.* 2004 (A)

The genus *Calosphaeria*, typified by *C. princeps*, includes 114 names while an asexual morph of *C. pulchella* was described in the monotypic genus *Calosphaeriophora* as *Ca. pulchella* (Réblová *et al.* 2004). *Calosphaeria pulchella* is a saprobic fungus occurring on various woody plants and has been recently isolated from wood of sweet cherry trees showing canker symptoms (Trouillas *et al.* 2010a, Berbegal *et al.* 2014). If *Calosphaeria princeps* and *C. pulchella* are congeneric, then these generic names are synonyms as suggested by Réblová *et al.* (2015). Based on priority, the number of species in the genus, and its widespread use, the generic name *Calosphaeria* is recommended for use.

Use *Pleurostoma* Tul. & C. Tul. 1863 (S) rather than *Pleurostomophora* Vijaykr. *et al.* 2004 (A)

The genus *Pleurostoma*, typified by *P. candollei*, includes six names, two of which have been placed elsewhere. One species, *P. ootheca*, has an asexual morph described as *Pleurostomophora ootheca*, the type species of *Pleurostomophora* (Vijaykrishna *et al.* 2004). Réblová *et al.* (2015) explored the phylogeny of this group including one species of *Pleurostoma* and four species of *Pleurostomophora*, although the type species of *Pleurostoma* was not available. While both sexual species are lignicolous, the asexual species were isolated from woody plants, soil or sewage (Schol-Schwarz 1970). They were also identified as etiological agents of subcutaneous phaeohyphomycosis (Meyer *et al.* 1975, Hironaga *et al.* 1989), and one asexual species of *Pleurostoma*, *P. ochraceum*, is known to cause human eumycetoma (Mhmoud *et al.* 2012). Assuming that *Pleurostoma candollei* and *P. ootheca* are congeneric, these generic names are synonyms. Given that the name *Pleurostoma* has priority, has the greater number of species, and has been widely used, we recommend the use of *Pleurostoma*. Based on these conclusions and recent molecular data, the three exclusively asexual *Pleurostomophora* species were transferred to *Pleurostoma* by Réblová *et al.* (2015).

Use *Togniniella* Réblová *et al.* 2004 (S) rather than *Phaeocrella* Réblová *et al.* 2004 (A)

The monotypic genera *Togniniella*, typified by *T. acerosa*, and *Phaeocrella*, typified by *P. acerosa*, were described in the same article for the sexual and asexual morphs of the same species and thus are synonyms having equal priority (Réblová *et al.* 2004). Although an older epithet was discovered for *T. acerosa* (Réblová 2011), this was later rescinded (Réblová *et al.* 2015). We recommend the use of the generic name *Togniniella*.

Chaetosphaeriales

Use *Chloridium* Link 1809 (A) rather than *Gonytrichum* Nees & T. Nees 1818 (A) or *Melanopsammella* Höhn. 1920 (S)

The type species of *Chloridium*, *C. viride* as *C. virescens*, has long been known to be a synonym of *Chaetosphaeria vermicularioides* (syn. *Melanopsammella vermicularioides*), a species that is congeneric with *M. inaequalis*, the type of *Melanopsammella* (Réblová & Winka 2000, Fernández *et al.*

2006, Crous *et al.* 2012). The type species of *Gonytrichum*, *G. caesium*, is the asexual morph of *M. inaequalis* (Hughes 1951), thus *Chloridium*, *Gonytrichum* and *Melanopsammella* are synonyms. Over 60 names have been placed in *Chloridium*, the most well-known of these three generic names. *Chloridium* was monographed by Gams & Holubová-Jechová (1976), however, many of these names have been removed to other genera. *Gonytrichum* includes over 20 species while *Melanopsammella* consists of five species, two of which already have names in *Chloridium* (Réblová *et al.* 1999, Réblová 2000). *Melanopsammella* has been distinguished from *Chaetosphaeria* by several authors (Réblová & Winka 2000, Fernández & Huhndorf 2005, Fernández *et al.* 2006, Crous *et al.* 2012). Given the widespread use of *Chloridium* and its priority, the generic name *Chloridium* is recommended for use.

New combinations:

Chloridium caesium* (Nees) Réblová & Seifert, *comb. nov.

MycoBank MB816825

Basionym: *Gonytrichum caesium* Nees, *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* **9**: 244 (1818).

Synonyms: *Eriosphaeria inaequalis* Grove, *J. Bot.*, Lond. **24**: 132 (1886).

Melanopsammella inaequalis (Grove) Höhn., *Ann. Mycol.* **17**: 121 (1920).

Chloridium chloroconium* (W. Gams & Hol.-Jech.) Réblová & Seifert, *comb. nov.

MycoBank MB816826

Basionym: *Chaetosphaeria chloroconia* W. Gams & Hol.-Jech., *Stud. Mycol.* **13**: 86 (1976).

Synonyms: *Melanopsammella chloroconia* (W. Gams & Hol.-Jech.) Réblová *et al.*, *Sydowia* **51**: 65 (1999).

Gonytrichum chlamydosporium G.L. Barron & G.C. Bhatt, *Mycopath. Mycol. appl.* **32**: 126 (1967).

Because the name *Chloridium chlamydosporium* (J.F.H. Beyma) S. Hughes 1958 already exists, and refers to a different taxon, the next older epithet for this species is taken up and placed in *Chloridium*.

Chloridium gonytrichii* (F.A. Fernández & Huhndorf) Réblová & Seifert, *comb. nov.

MycoBank MB816827

Basionym: *Melanopsammella gonytrichii* F.A. Fernández & Huhndorf, *Fungal Diversity* **18**: 42 (2005).

Crous *et al.* (2012) showed that this species is closely related to *Chloridium virescens* as *Melanopsammella vermicularioides*.

Use *Menispora* Pers. (A) 1822 rather than *Zignoëlla* Sacc. 1878 (S)

The genus *Menispora* is typified by *M. glauca*, which has a sexual morph referred to as *Chaetosphaeria glauca* (Holubová-Jechová 1973a), which is a synonym of the earlier name *Zignoëlla ovoidea*. The type of *Zignoëlla* is *Z. pulviscula*, which has an asexual morph known as *Menispora*

caesia (Booth 1957, Constantinescu *et al.* 1995). Fernández *et al.* (2006) demonstrated that *Zignoëlla* is distinct from *Chaetosphaeria* based on *C. innumera* and that *Z. pulviscula* and *Z. ovoidea* are congeneric, thus *Menispora* and *Zignoëlla* are synonyms. Like *Chaetosphaeria*, *Zignoëlla* has been used as a repository for species that have a similar morphology, namely black non-stromatic ascomata with a papilla and hyaline, septate ascospores, but are phylogenetically diverse. Given the lack of characterization of *Zignoëlla*, the greater use of *Menispora*, and following the principle of priority, *Menispora* is recommended for use.

Use *Menisporopsis* S. Hughes 1952 (A) rather than *Menisporopascus* Matsush. 2003 (S)

The monotype species of *Menisporopascus*, *M. kobensis*, was described for the asexual morph *Menisporopsis kobensis*. Because *M. kobensis* is congeneric with the type species of *Menisporopsis*, *M. theobromae*, as indicated by Rodrigues de Cruz *et al.* (2014), *Menisporopsis* and *Menisporopascus* are synonyms. *Menisporopsis* includes 10 names, is widely known, and has priority, therefore, the use of *Menisporopsis* is recommended.

Use *Sporoschisma* Berk. & Broome 1847 (A) rather than *Melanochaeta* E. Müll. *et al.* 1969 (S)

The genus *Sporoschisma*, typified by *S. mirabile*, is considered the asexual morph of *Melanochaeta aotearoae*, while the type of *Melanochaeta*, *M. hemipsila*, is the sexual morph of *S. saccardoi*, now considered a synonym of *S. hemipsila* (Sivichai *et al.* 2000, Zelski *et al.* 2014a). *Sporoschisma mirabile* and *S. hemipsila* were shown to be congeneric (Fernández *et al.* 2006), thus *Sporoschisma* and *Melanochaeta* are synonyms. Of the five species in *Melanochaeta*, four have *Sporoschisma* asexual morphs (Sivichai *et al.* 2000, Mugambi & Huhndorf 2008). *Sporoschisma* includes 23 names, some of which have been placed elsewhere. If *Melanochaeta* were used, many name changes would be required. Based on priority and fewer number of name changes, we follow Zelski *et al.* (2014a) and recommend the use of *Sporoschisma* for this genus.

New combinations:

Sporoschisma daemonoropis* (J. Fröhl. & K.D. Hyde) A.N. Mill., *comb. nov.

MycoBank MB816828

Basionym: *Melanochaeta daemonoropis* J. Fröhl. & K.D. Hyde, *Palm Microfungi*: 235 (2000).

Sporoschisma taitense* (Mugambi & Huhndorf) A.N. Mill., *comb. nov.

MycoBank MB816829

Basionym: *Melanochaeta taitensis* Mugambi & Huhndorf, *Sydowia* **60**: 263 (2008).

A *Sporoschisma* asexual morph was noted to occur with this type specimen.

Use *Stanjehughesia* Subram. 1992 (A) rather than *Umbrinosphaeria* Réblová 1999 (S)

Stanjehughesia hormiscioides, type of *Stanjehughesia*, was established with cultivation techniques as an asexual morph of *Umbrinosphaeria caesariata*, the monotype species of *Umbrinosphaeria* (Réblová 1999), therefore *Stanjehughesia* and *Umbrinosphaeria* are synonyms. In exploring species of *Sporidesmium*, Shenoy *et al.* (2006) demonstrated the polyphyletic nature of *Stanjehughesia*; however, 15 species have been described in that genus and the name is well-known. Given its priority, we recommend the use of *Stanjehughesia*.

Coniochaetales

Use *Coniochaeta* (Sacc.) Cooke 1887 (S) rather than *Lecytophora* Nannf. 1934 (A)

The genus *Coniochaeta*, typified by *C. ligniaria*, was shown to be congeneric with *Lecytophora*, typified by *L. lignicola* (Weber *et al.* 2002, Damm *et al.* 2010, Khan *et al.* 2013, Perdomo *et al.* 2013). *Coniochaeta* is a well-known genus with almost 100 names while *Lecytophora* includes nine names, two of which are linked to species of *Coniochaeta*. These fungi are known as endophytes, pathogens of woody trees, saprobes in terrestrial and freshwater habitats (Raja *et al.* 2012) and, to a lesser degree, human pathogens (Damm *et al.* 2010). Both Khan *et al.* (2013) and Perdomo *et al.* (2013) include species of *Lecytophora* in a clade with the type species of *Coniochaeta*. Given the number of species, its widespread use, following the principle of priority, and in agreement with Khan *et al.* (2013), *Coniochaeta* is recommended for use.

Conioscyphales

Use *Conioscypha* Höhn. 1904 (A) rather than *Conioscyphascus* Réblová & Seifert 2004 (S)

The genus *Conioscypha*, with *C. lignicola* as type species, includes 13 species from freshwater and terrestrial habitats. The genus *Conioscyphascus*, typified by *Ca. varius*, was originally established for fungi with *Conioscypha* asexual morphs (Réblová & Seifert 2004a). The two sexual species, *Ca. gracilis* with *C. gracilis* as its asexual morph and *Ca. varius* with *C. varia*, were recently placed in *Conioscypha* (Zelski *et al.* 2014b, Réblová *et al.* 2016). Although *C. lignicola* and 10 other species of *Conioscypha* are known only in their asexual morph, using DNA sequence data and cultivation studies, the link between sexual and asexual morphs has been experimentally proven for *C. peruviana* (Zelski *et al.* 2014b) and *C. varia* (Réblová & Seifert 2004a). Such a link has not yet been confirmed for the third species known to have a sexual morph, *C. gracilis*, however, conidia were observed on the substratum near ascomata (Réblová & Seifert 2004a, Zelski *et al.* 2014b). The molecular data clearly show that *Conioscypha* and *Conioscyphascus* are congeneric (Réblová & Seifert 2004a). Based on priority and the greater number of species, the name *Conioscypha* is recommended for use.

Cordanales

Use *Cordana* Preuss 1851 (A) rather than *Pseudobotrytis* Krzemien. & Badura 1954 (A) or *Porosphaerella* E. Müll. & Samuels 1982 (S)

The type species of *Cordana*, *C. pauciseptata*, was linked to a sexual morph described as *Porosphaerella cordanophora*, type of *Porosphaerella* (Müller & Samuels 1982), thus these generic names are synonyms. *Cordana* is a well-known asexual genus that includes 22 names while three species have been placed in *Porosphaerella*. Réblová and Seifert (2007) showed that *P. borinquensis* and *P. cordanophora* are closely related despite the fact that *P. borinquensis* has an asexual morph regarded as *Pseudobotrytis terrestris* which Hernández-Restrepo *et al.* (2014) recognized as *Cordana terrestris* including *Porosphaerella borinquensis* as a synonym. They also considered the type species of *Pseudobotrytis*, *P. fusca*, to be a synonym of *C. terrestris*, thus this generic name with four species is also a synonym of *Cordana*. Given its priority, fewer number of name changes required, and widespread use, we recommend the use of *Cordana* rather than *Pseudobotrytis* or *Porosphaerella*.

Coronophorales

Protect *Chaetosphaerella* E. Müll. & C. Booth 1972 (S) over *Oedemium* Link 1824 (A) and rather than *Veramycina* Subram. 1995 (A)

The genus *Chaetosphaerella*, typified by *C. phaeostroma*, has an asexual morph regarded as *Oedemium minus* (Booth 1958, Réblová 1999). The type species of *Oedemium*, *O. atrum* as *O. didymium*, is the asexual morph of *Chaetosphaerella fusca* (Ellis 1971). These two species of *Chaetosphaerella* were shown to be congeneric (Mugambi & Huhndorf 2010), thus *Chaetosphaerella* and *Oedemium* are synonyms. *Veramycina elegans*, the monotype species of *Veramycina*, was regarded as a later name for the asexual morph of *C. phaeostroma* (Réblová 1999), thus this generic name is also a synonym of *Chaetosphaerella*. Because *Veramycina* has been rarely used, that name is not a contender for use, however, the decision of whether to use *Oedemium* or *Chaetosphaerella* is more difficult. The genus *Oedemium* was clarified and monographed including two species by Hughes & Hennebert (1963) both of which have been linked with *Chaetosphaerella*. Although more species have been described in *Oedemium*, some have been placed elsewhere and others remain obscure. *Chaetosphaerella* has been placed phylogenetically in *Chaetosphaerellaceae*, *Coronophorales* (Mugambi & Huhndorf 2010). In the past decade *Chaetosphaerella* appears in the literature much more frequently than *Oedemium*, thus we recommend *Chaetosphaerella* for protection.

Glomerellales

Use *Colletotrichum* Corda 1837 (A) rather than *Glomerella* Spauld. & Schrenk 1903 (S)

The genus *Colletotrichum*, typified by *C. lineola*, is a large genus with 817 epithets (*Index Fungorum* 2016), many of which are considered plant pathogens. The number of

accepted species has varied over the decades with most species previously considered to be host-specific. Based on von Arx (1957) in which several hundred names were placed in synonymy with *C. gloeosporioides*, the number of species in *Colletotrichum* decreased to 11. Sutton (1980) included about 22 species in *Colletotrichum* for which he provided a key. With the use of molecular sequence data, the number of recognized species has increased with some species determined to be host-specific while others have a broad host range (Rojas et al. 2010, Damm et al. 2009, 2012, Weir et al. 2012). The type species, *Colletotrichum lineola*, had long been considered to be a synonym of *C. dematium* but was recently recognized as a distinct species in the *C. dematium* species complex (Damm et al. 2009, Cannon et al. 2012). Meanwhile, the sexual morphs of *Colletotrichum* were placed in the genus *Glomerella*, typified by *G. cingulata*, which has been considered the sexual morph of *C. gloeosporioides*. Recently this relationship has been questioned with *C. gloeosporioides* and *G. cingulata* regarded as distinct species in the *C. gloeosporioides* species complex (Weir et al. 2012). Over 100 names have been described in *Glomerella*, many of which have names in *Colletotrichum*. Although considered different species, there is no question that *Colletotrichum* and *Glomerella* represent the same genus.

Use of *Colletotrichum* over *Glomerella* is recommended for use for several reasons. Species of *Colletotrichum* are mainly associated with plants as endophytes and pathogens and have been thoroughly studied. In contrast, their sexual morphs regarded as *Glomerella* tend to occur on dead plant material and are less frequently encountered (Cannon et al. 2012, Maharachchikumbura et al. 2015). *Colletotrichum* is the older generic name and is well established in the plant pathological and plant breeder literature (Cannon et al. 2012). Preference has been given to *Colletotrichum* by several groups working on this genus (Cannon et al. 2012, Maharachchikumbura et al. 2015) based on its widespread use, greater number of names, and priority.

Use *Cylindrotrichum* Bonord. 1851 (A) rather than *Reticulascus* Réblová & W. Gams 2011 (S)

The type species of *Cylindrotrichum*, *C. oligospermum*, was shown to represent the asexual morph of *Chaetosphaeria tulasneorum*, the type species of *Reticulascus* (Réblová & Gams 1999, Réblová et al. 2011a), thus the generic names *Cylindrotrichum* and *Reticulascus* are synonyms. *Cylindrotrichum* includes 23 names (Rambelli & Onofri 1987) while *Reticulascus* includes only two names. Between these two generic names, *Cylindrotrichum* is the most widely known. Given its widespread use, priority, and greater number of names, we recommend the use of *Cylindrotrichum* rather than *Reticulascus*.

Use *Monilochaetes* Halst. ex Harter 1916 (A) rather than *Australiasca* Sivan. & Alcorn 2002 (S)

The type species of *Monilochaetes*, *M. infuscans*, was shown to be congeneric with *Australiasca queenslandica*, the type species of *Australiasca* (Réblová et al. 2011a). The two species described in *Australiasca* have names in *Monilochaetes*, thus, if the older name *Monilochaetes* is used, no name changes are required. This generic name

includes six species and has priority, thus *Monilochaetes* is recommended for use.

Use *Plectosphaerella* Kleb. 1929 (S) rather than *Plectosporium* M.E. Palm et al. 1995 (A)

The genus *Plectosporium*, typified by *P. tabacinum*, was established for the asexual morph of species of *Plectosphaerella* of which the type species, *P. cucumeris*, is the name applied to the sexual morph of *Plectosporium tabacinum* (Palm et al. 1995), thus *Plectosphaerella* and *Plectosporium* are synonyms. *Plectosphaerella cucumerinum* is known as the cause of fruit and collar rot, and collapse of several crops including melons (Carlucci et al. 2012). *Plectosphaerella* includes 14 names while four species have been placed in *Plectosporium*. Carlucci et al. (2012) provided a molecular account of *Plectosphaerella* in the *Plectosphaerellaceae* and transferred species of *Plectosporium* to *Plectosphaerella*. Given its priority, greater number of names, economic importance, and recent account of the genus, the use of *Plectosphaerella* is recommended.

Use *Sporoschismopsis* Hol.-Jech. & Hennebert 1972 (A) rather than *Porosphaerellopsis* Samuels & E. Müll. 1982 (S)

The generic name *Sporoschismopsis* was introduced by Holubová-Jechová & Hennebert (1972) with *S. moravica* as type species and it now includes eight names (Holubová-Jechová 1973b). The generic name *Porosphaerellopsis*, typified by *P. sporoschismophora* (Samuels & Müller 1978, Müller & Samuels 1982), was established for the sexual morphs of *Sporoschismopsis*. *Porosphaerellopsis sporoschismophora* was transferred to *Sporoschismopsis* (Réblová 2014) while a second species, *P. bipolaris* described by Ranghoo et al. (2001), was excluded from the genus. Another sexual morph was experimentally proven for *S. angustata* (Réblová 2014). Based on DNA sequence data, the two asexual morphs, *S. angustata* and *S. sporoschismophora*, form a strongly supported monophyletic clade in the *Reticulascaceae*, *Glomerellales*. Given that *S. angustata*, *S. moravica*, and *S. sporoschismophora* are congeneric, the generic names *Sporoschismopsis* and *Porosphaerellopsis* are synonyms. Based on priority and the greater number of species, *Sporoschismopsis* is recommended for use.

Halosphaeriales

Use *Corollospora* Werderm. 1922 (S) rather than *Varicosporina* Meyers & Kohlm. 1965 (A) and *Halosigma* Nakagiri et al. 2009 (A)

Although the type species of *Corollospora*, *C. maritima*, does not appear to have an asexual morph in *Varicosporina*, a number of species of *Corollospora* are linked to varicosporina-like asexual morphs. Abdel-Wahab et al. (2009) showed that the type of *Varicosporina*, *V. ramulosa*, groups within *Corollospora*, thus *Corollospora* and *Varicosporina* are synonyms. The type species of *Halosigma*, *H. luteola* as *Sigma* *luteola*, was shown by culture techniques to be the asexual morph of *Corollospora luteola* (Nakagiri & Tokura, 1982). Jones et al. (2009) demonstrated that the three marine species

recognized in *Halosigmoidea* grouped with *Corollospora*, thus *Halosigmoidea* is also a synonym of *Corollospora*. Over 20 species are included in *Corollospora* while only three names have been described in *Varicosporina*, two of which have names in *Corollospora*, and only three names were included in *Halosigmoidea*. Given the number of names, widespread use, and priority, we recommend the use of *Corollospora*.

New combinations:

Corollospora marina* (Haythorn & E.B.G. Jones) E.B.G. Jones, K.L. Pang & Abdel-Wahab, *comb. nov.

MycoBank MB816830

Basionym: *Sigmoidea marina* Haythorn & E.B.G. Jones, *Trans. Brit. Mycol. Soc.* **74**: 620 (1980).

Synonym: *Halosigmoidea marina* (Haythorn & E.B.G. Jones) Nakagiri *et al.*, *Bot. Mar.* **52**: 355 (2009).

Corollospora parvula* (Zuccaro *et al.*) E.B.G. Jones, K.L. Pang & Abdel-Wahab, *comb. nov.

MycoBank MB816831

Basionym: *Halosigmoidea parvula* Zuccaro *et al.*, *Bot. Mar.* **52**: 355 (2009).

Corollospora ramulosa* (Meyers & Kohlm.) E.B.G. Jones & Abdel-Wahab, *comb. nov.

MycoBank MB816832

Basionym: *Varicosporina ramulosa* Meyers & Kohlm., *Canad. J. Bot.* **43**: 916 (1965).

Melanosporales

Use *Melanospora* Corda 1836 (S) rather than *Gonatobotrys* Corda 1839 (A)

The type species of *Melanospora*, *M. zamiae*, has been shown to be congeneric with *M. damnosa* (Cannon & Hawksworth 1982), of which the type species of *Gonatobotrys*, *G. simplex*, is considered a synonym (Vakili 1989). Given the widespread use of *Melanospora*, its priority, and the 120 names in *Melanospora* while *Gonatobotrys* includes only 32 names, *Melanospora* is recommended for use.

New combination:

Melanospora simplex* (Corda) D. Hawksw., *comb. nov.

MycoBank MB816833

Basionym: *Gonatobotrys simplex* Corda, *Prachtflora*: 9 (1839).

Synonyms: *Sphaeroderma damnosum* Sacc. & Berl., *Riv. Patol. Veg.* **4**: 56 (1896) [“1895”].

Melanospora damnosa (Sacc.) Lindau, *Nat. Pflanzenfam.* **1** (1*): 353 (1897).

Microascales

Use *Knoxdaviesia* M.J. Wingf. *et al.* 1988 (A) rather than *Gondwanamyces* G.J. Marais & M.J. Wingf. 1998 (S)

Knoxdaviesia proteae, type of the genus *Knoxdaviesia*, was described as the asexual morph of *Ceratocystiopsis proteae*, a species that was later placed in the genus *Gondwanamyces* as the type species, thus *Knoxdaviesia* and *Gondwanamyces* are synonyms. Both generic names have been used about equally. In agreement with de Beer *et al.* (2013), the use of the oldest name, *Knoxdaviesia*, is recommended.

Use *Sphaeronaemella* P. Karst. 1884 (S) rather than *Gabarnaudia* Samson & W. Gams 1974 (A)

The type species of *Sphaeronaemella*, *S. helvellae*, is congeneric with the type species of *Gabarnaudia*, *G. betae*, described as the asexual morph of *S. fimicola* (Samson 1974). Confirmed by molecular data (Hausner & Reid 2004), these generic names are synonyms. Five species of *Gabarnaudia* have been placed in *Sphaeronaemella* (De Beer *et al.* 2013), a genus with 30 species. *Sphaeronaemella* is most commonly used and has priority, thus we recommend the use of *Sphaeronaemella*.

Phyllachorales

Use *Diachora* Müll. Arg. 1893 (S) rather than *Diachorella* Höhn. 1918 (A)

The generic names *Diachora*, typified by *D. onobrychidis*, and *Diachorella*, typified by *D. onobrychidis*, are apparently based on the same type specimen, thus these names are synonyms. *Diachora* is considered a name for the sexual morph while *Diachorella* represents the asexual morph. Both generic names include less than ten species. Müller (1986) and Cannon (1991) reviewed species in these genera including the type species. Because *Diachora* is more widely known and has priority, we recommend that name for use.

Use *Ophiodothella* (Henn.) Höhn. 1904 (S) rather than *Acerviclypeata* Hanlin 1990 (A)

The genus *Ophiodothella*, typified by *O. atromaculans*, now includes 26 species, which are obligate parasites of subtropical and tropical plants (Hanlin *et al.* 1992). An asexual morph of *O. vaccinii*, now *O. angustissima*, cause of flyspeck leafspot of *Vaccinium arboretum* in the southeastern United States, was described as the monotype species of *Acerviclypeatus*, *A. poriformans* (Hanlin 1990), thus *Ophiodothella* and *Acerviclypeatus* are synonyms. Not much is known about *O. atromaculans* but this species appears to be congeneric with *O. angustissima* (Hanlin & González 2013). Given the number of species and priority, we recommend the use of *Ophiodothella*.

Use *Polystigma* DC 1815 (S) rather than *Polystigmina* Sacc. 1884 (A) and *Rhodoseptoria* Naumov 1913 (A)

The generic names *Polystigma*, typified by *P. rubrum*, and *Polystigmina*, typified by *P. rubrum*, are based on the same basionym of their respective type species and, therefore, are synonyms for the sexual and asexual morphs. Cannon (1996) monographed *Polystigma* including five species on *Prunus* among others. He also noted that the type species of the monotypic *Rhodoseptoria* was a synonym of *Polystigma*

rubra, thus that generic name is also a synonym of *Polystigma*. Several species of *Polystigma* cause diseases such as red blotch of almonds caused by *P. amygdalinum* and red leaf spot of plum caused by *P. rubrum* (Habibi et al. 2015). All but two of the seven taxa placed in *Polystigmata* are synonyms of *Polystigma rubra* while *Polystigma* includes over 50 species. Given the greater number of species, its widespread use, economic importance, and priority, we recommend the use of *Polystigma*.

Use *Pseudothiella* Petr. 1928 (S) rather than *Pseudothiopsella* Petr. 1928 (A)

The monotypic generic names *Pseudothiella*, typified by *P. hirtellae*, and *Pseudothiopsella*, typified by *P. hirtellae*, were described at the same time for sexual and asexual morphs of the same species, thus they are synonyms with equal priority. *Pseudothiella* has been used slightly more often than *Pseudothiopsella* (Furlanetto & Dianese 1998), therefore we recommend use of *Pseudothiella*.

Pleurotheciales

Use *Pleurothecium* Höhn. 1919 (A) rather than *Carpoligna* F.A. Fernández & Huhndorf 1999 (S)

The genus *Pleurothecium* is typified by *P. recurvatum*. A sexual morph is known for *P. recurvatum* for which the monotypic generic name *Carpoligna*, typified by *C. pleurothecii*, was introduced (Fernández et al. 1999). Of the eight species assigned to the genus *Pleurothecium*, only three have DNA sequence data. *Pleurothecium recurvatum* and *P. semifecundum* form a strongly supported monophyletic clade in the *Pleurotheciales* and represent the core of the genus (Réblová et al. 2016), while *P. obovoideum* (Arzanlou et al. 2007) is nested in the same order but within another clade. Given its priority and greater number of species, *Pleurothecium* is recommended for use.

Savoryellales

Use *Canalisporium* Nawawi & Kuthub. 1989 (A) rather than *Ascothailandia* Sri-indr. et al. 2010 (S)

The genus *Canalisporium*, typified by *C. caribense*, includes 12 species (Nawawi & Kuthubuteen 1989). A sexual morph was described for *C. grenadoideum* and linked to the monotypic genus *Ascothailandia*, typified by *A. grenadoidea* (Sri-indrasudthi et al. 2010). The link between sexual and asexual morphs of this species was confirmed experimentally in axenic culture. Six species of *Canalisporium*, including the type species, were the subject of phylogenetic analysis based on DNA sequence data of three nuclear loci by Sri-indrasudthi et al. (2010) who confirmed that *Ascothailandia* and *Canalisporium* are congeneric. Based on priority and the greater number of species, the generic name *Canalisporium* is recommended for use.

Sordariales

Use *Chaetomium* Kunze 1817 (S) rather than *Trichocladium* Harz 1871 (A), *Botryotrichum* Sacc. & Marchal 1885 (A), or *Humicola* Traaen 1914 (A)

Chaetomium, typified by *C. globosum*, is a well-known genus of cellulose-decomposing fungi some of which cause diseases of humans. The sexual morph of *Botryotrichum piluliferum*, type of the asexual genus *Botryotrichum*, was described as *Chaetomium piluliferum* (Daniels 1961). Later this species was confirmed to be congeneric with *C. globosum* (Untereiner et al. 2001, Nonaka et al. 2012), thus *Chaetomium* and *Botryotrichum* are synonyms. Although sexual states have not been discovered for the type of *Humicola*, *H. fuscoatra* or related species such as the common *H. grisea*, nuc28S rDNA sequences place these species within the present concept of *Chaetomium*. The type of *Trichocladium*, *T. asperum*, also lacks a sexual morph and occurs in the same clade (Hambleton et al. 2005). Over 500 names have been described in *Chaetomium*. *Botryotrichum* currently includes eight species. *Humicola* and *Trichocladium* each include about 20 species, but both are phylogenetically heterogeneous and most species belong to different clades. Given its extensive use and economic importance, its priority, and the greater number of species, we recommend the use of *Chaetomium*.

Use *Lasiosphaeris* Clem. 1909 (S) rather than *Lasiadelphia* Réblová & W. Gams 2011 (A)

The genus *Lasiosphaeris* was described by Clements (1909) with *L. hispida* as its type species. The name was reestablished by Miller & Huhndorf (2004) for species segregated from *Lasiosphaeria* based on nuc28S rDNA sequences. Two species were accepted in the genus, *L. hispida* and *L. hirsuta*. Gams & Holubová-Jechová (1976) introduced *Phialophora* sect. *Catenulatae*, typified by the dematiaceous species *Phialophora lasiosphaeriae*, the asexual morph experimentally proven for *L. hispida* (as *L. hirsuta*, Gams & Holubová-Jechová 1976, Gams 2000). Réblová et al. (2011b) described the generic name *Lasiadelphia* with *L. lasiosphaeriae* as the type species for the asexual morph of *Lasiosphaeris hispida*. Therefore, the type species of *Lasiosphaeris* and *Lasiadelphia* are morphs of the same species and the generic names are synonyms. Based on priority and the greater number of species, the generic name *Lasiosphaeris* should be used.

Use *Mammaria* Rabenh. 1854 (A) rather than *Pseudocercophora* Subram. & Sekar 1986 (S)

The genus *Mammaria*, typified by *M. echinobotryoides*, includes two names while the monotypic genus *Pseudocercophora*, typified by *P. ingoldii*, was described for the sexual morph of *M. echinobotryoides*, thus these generic names are synonyms. If the principle of priority is followed, no names changes are required, thus, the use of *Mammaria* is recommended.

Use *Neurospora* Shear & B.O. Dodge 1927 (S) rather than *Chrysonilia* Arx 1981 (A)

The genus *Neurospora*, typified by *N. sitophila*, includes the well-known model organism, *N. crassa*. The asexual

morph of the type species was described in *Chrysonilia* with *C. sitophila* as the type species. There is no question that *Neurospora* and *Chrysonilia* are generic synonyms (von Arx 1981). At present *Neurospora* includes 57 names while only three names have been placed in *Chrysonilia*, all of which also have names in *Neurospora*. Given its priority, the number of species, and its widespread use, it is recommended that *Neurospora* be used rather than *Chrysonilia*.

Use *Stromatographium* Henn. 1907 (A) rather than *Fluviostroma* Samuels & E. Müll. 1980 (S)

The genus *Stromatographium*, typified by *S. stromaticum*, is the asexual state of *Fluviostroma wrightii*, monotype species of *Fluviostroma* (Samuels & Müller 1980). Seifert (1987) clarified the confusion between *Stromatographium* and *Stromatostilbella* and showed that the asexual morph of *F. wrightii* belongs in *Stromatographium*. Given its greater use and priority, we recommend the use of *Stromatographium*.

Togniniales

Protect *Phaeoacremonium* W. Gams 1996 (A) over *Togninia* Berl. 1900 (S) and *Romellia* Berl. 1900 (S)

The genus *Phaeoacremonium* was established for *P. parasiticum*, first described causing phaeohyphomycosis in humans, but also causing wilts in tropical trees and an economically important disease of grapevine known as esca disease (Hawksworth *et al.* 1976, Crous *et al.* 1996). Once this genus was described, a number of species were discovered and placed in *Phaeoacremonium*. Based on phylogenetic analysis and mating studies, sexual morphs were found for these species in *Togninia*, formerly attributed to *Calosphaeriales* (Mostert *et al.* 2003). Species of *Phaeoacremonium* are known as vascular plant pathogens such as esca and Petri disease of grapevine as well as human pathogens. Two monographic accounts of *Phaeoacremonium* have been published (Mostert *et al.* 2006, Réblová *et al.* 2015). Gramaje *et al.* (2015) formally proposed to protect the name *Phaeoacremonium* against *Togninia* because although *Togninia* has 26 epithets, half have not been reported since their description, and only nine species have been experimentally linked with *Phaeoacremonium*. *Phaeoacremonium* includes 46 species of which all are known from culture and their DNA data are available. Also *Phaeoacremonium* is favoured by plant pathologists and medical mycologists and is well established in the literature. All required new combinations were made in Gramaje *et al.* (2015). The type species of *Romellia*, *R. vibratilis*, was sequenced and placed in *Togninia* by Réblová & Mostert (2007) and recently moved to *Phaeoacremonium* by Gramaje *et al.* (2015). Three of the four species of *Romellia* have since been placed elsewhere. Thus, because it is widely used and has the greatest number of species, we recommend the protection of *Phaeoacremonium* over *Togninia* and *Romellia*.

Torpedosporales

Use *Juncigena* Kohlm. *et al.* 1997 (S) rather than *Moheitospora* Abdel-Wahab *et al.* 2010 (A)

The monotypic genus *Juncigena* is typified by *J. adarca* for which an asexual morph was described as *Cirrenalia adarca* (Kohlmeyer *et al.* 1997). Later the asexual species was placed in the genus *Moheitospora* as *M. adarca* along with the type species, *M. fruticosae* (Abdel-Wahab *et al.* 2010), thus *Juncigena* and *Moheitospora* are synonyms. Although one name change would be required, *Juncigena* has priority and has greater use in the literature, thus it is recommended that *Juncigena* be used. Schoch *et al.* (2007) showed that this genus belongs in *Hypocreomycetidae* and later Jones *et al.* (2015) placed it in *Juncigenaceae*, *Torpedosporales*.

New combination:

Juncigena fruticosae* (Abdel-Wahab *et al.*) A.N. Mill. & Shearer, *comb. nov.

Mycobank MB816834

Basionym: *Moheitospora fruticosae* Abdel-Wahab *et al.*, *Mycol. Progr.* 9: 551 (2010).

Use *Torpedospora* Meyers 1957 (S) rather than *Glomerulispora* Abdel-Wahab & Nagah. 2010 (A)

The monotype species of *Glomerulispora*, *G. mangrovei*, groups with the two species of *Torpedospora* such that these generic names are considered synonyms (Abdel-Wahab *et al.* 2010, Jones *et al.* 2015). These species are marine fungi found on driftwood throughout the world. Given the widespread use of *Torpedospora* and its priority, use of *Torpedospora* is recommended.

New combination:

Torpedospora mangrovei* (Abdel-Wahab & Nagah.) E.B.G. Jones & Abdel-Wahab, *comb. nov.

Mycobank MB816835

Basionym: *Glomerulispora mangrovei* Abdel-Wahab & Nagah., *Mycol. Progr.* 9: 553 (2010).

Xylariales

Use *Arthrimum* Kunze 1817 (A) rather than *Apiospora* Sacc. 1875 (S), *Pteroconium* Sacc. ex Grove 1914 (A), and *Scyphospora* L.A. Kantsch 1928 (A)

Species in the genus *Arthrimum* are widespread and commonly encountered as saprobes and secondary invaders especially on monocotyledonous plants. The common species, *A. arundinis*, is the name typified by the sexual morph of the type species of *Apiospora*, *A. montagnei* (Müller & Arx 1962). Less is known about the type species of *Arthrimum*, *A. caricicola*, but this species is assumed to be congeneric with *A. arundinis* (Crous & Groenewald 2013), although they were unable to find material to include in their molecular phylogeny. Crous & Groenewald (2013) also showed that *Pteroconium asteroides*, type of *Pteroconium*,

is a synonym of *Arthrimum*. Samuels et al. (1981) suggested that *Scyphospora phyllostachydis* was the asexual morph of *Apiospora tintinnabula*, now *Arthrimum hysterinum*, thus *Scyphospora* is a synonym of *Arthrimum*. In agreement with both Crous & Groenewald (2013) and Senanayake et al. (2015), *Arthrimum* has priority and is the most widely used of these generic names, thus use of *Arthrimum* is recommended.

Although Hughes (1958) followed by Ellis (1965), Samuels et al. (1981) and Crous & Groenewald (2013) suggested that *Cordella* is a synonym of *Arthrimum*, this is based on the assumption that *C. coniosporioides* is the type species. *Cordella* was described with two species without indicating a type. Clements & Shear (1931) were the first to select a type species for *Cordella* and they list *C. spinulosa*, a species that Hughes (1958) placed in *Melanographium*, thus *Cordella* is not a synonym of *Arthrimum*.

Based on the molecular sequence data provided by Crous & Groenewald (2013), *Apiospora sinensis* is placed in *Arthrimum*.

New combination:

Arthrimum sinense (K.D. Hyde et al.) Crous & J.Z. Groenew., **comb. nov.**

MycoBank MB816836

Basionym: *Apiospora sinensis* K.D. Hyde et al., *Sydowia* **50**: 27 (1998).

In addition a nomenclator is provided for *Arthrimum hysterinum* because of confusion about the correct name for this species and its numerous synonyms. Previously recognized in *Apiospora* by Tang et al. (2007), Kirk (1986) and Sivanesan (1983), Crous & Groenewald (2013) and Senanayake et al. (2015) included *Apiospora bambusae*, *A. setosa* and *A. tintinnabula* as synonyms within the genus *Arthrimum*. The correct name for this species is *Arthrimum hysterinum* with the synonyms listed below:

Arthrimum hysterinum (Sacc.) P.M. Kirk, *Trans. Brit. Mycol. Soc.* **86**: 409 (1986).

Basionym: *Melanconium hysterinum* Sacc., *Bolm Soc. broteriana*, *Coimbra*, sér. 1 **11**: 21 (1893).

Synonyms: *Scyphospora hysterina* (Sacc.) Sivan., *Trans. Brit. Mycol. Soc.* **81**: 331 (1983).

Melanconium bambusae Turconi, *Atti Ist. bot. R. Univ. Pavia*, sér.2 **16**: 251 (1916).

Scirrhia bambusae Turconi, *Atti Ist. bot. R. Univ. Pavia*, sér. 2 **16**: 531 (1916).

Scirrhodithis bambusae (Turconi) Trotter, in Saccardo, *Syll. Fung.* **24**: 611 (1926).

Placostroma bambusae (Turconi) R. Sprague, *Diseases Cereals Grasses N. Amer.*: 121 (1950).

Apiospora bambusae (Turconi) Sivan., *Trans. Brit. Mycol. Soc.* **81**: 331 (1983).

Scyphospora phyllostachydis L.A. Kantsch., *Boléz. Rast.* **17**: 88 (1928).

Cordella johnstonii M.B. Ellis, *Mycol. Pap.* **103**: 31 (1965).

Apiospora setosa Samuels et al., *New Zealand J. Bot.* **19**: 142 (1981).

Apiospora tintinnabula Samuels et al., *New Zealand J. Bot.* **19**: 142 (1981).

Use *Collodiscula* I. Hino & Katum. 1955 (S) rather than *Acanthodochium* Samuels et al. 1987 (A)

The genus *Collodiscula*, typified by *C. japonica*, was linked to *Acanthodochium collodisculae*, type of *Acanthodochium*, by Samuels et al. (1987), who considered these names morphs of the same species, thus these generic names are synonyms. Three species have been added to *Collodiscula* (Li et al. 2015a, b). Given that this generic name has priority and includes four species, the use of *Collodiscula* is recommended.

Use *Daldinia* Ces. & De Not. 1863 (S) rather than *Annelosporium* M.L. Davey 2010 (A) or *Versiumyces* Whalley & Watling 1989 (S)

The genus *Daldinia*, typified by *D. concentrica*, has recently been monographed by Stadler et al. (2014). They showed that the monotype species of *Annelosporium*, *A. nemorosa*, belongs in *Daldinia* as *D. nemorosa*, thus *Daldinia* and *Annelosporium* are synonyms. They also showed that the monotype species of *Versiumyces*, *V. cahuchucosus*, belongs in *Daldinia* as *D. cahuchucosa*, thus *Versiumyces* is a synonym of *Daldinia*. *Daldinia* is a well-known genus with many species and has priority, thus it is recommended for use.

Protect *Diatrype* Fr. 1849 (S) over *Libertella* Desm. 1830 (A)

The genus *Diatrype*, typified by *D. disciformis*, includes over 500 names (*Index Fungorum* 2016), some of which cause canker diseases of hardwoods such as *D. stigma* associated with diseased grapevines (Rolshausen et al. 2006). *Libertella betulina*, type of the genus *Libertella*, is known to be the asexual morph of *Diatrype stigma* (Grove 1937, Kutorga et al. 2006) and *D. disciformis* has an asexual morph described as *L. disciformis*. Trouillas et al. (2010b) showed that *D. disciformis* and *D. stigma* are congeneric, thus *Diatrype* and *Libertella* are synonyms. Because *Diatrype* is widely known, has a greater number of species, and includes important plant pathogenic fungi, *Diatrype* is recommended for protection over *Libertella*.

Use *Hypocreodendron* Henn. 1897 (A) rather than *Discoxylaria* J.C. Lindq. & J.E. Wright 1964 (S)

Hypocreodendron is a monotypic genus based upon *H. sanguineum* for which Lindqvist & Wright (1964) introduced the monotypic genus *Discoxylaria*, typified by *D. myrmecophila*, as the sexual morph. Rogers et al. (1995) observed a specimen of *D. myrmecophila* that bore both conidial and mature perithecial morphs. They compared their collection with the holotype specimen and obtained the conidial morph in culture initiated from ascospores to demonstrate that these were alternate morphs of the same species. Thus, *Hypocreodendron* and *Discoxylaria* are synonyms. In agreement with Stadler et al. (2013), Maharachchikumbura et al. (2015), and priority, we recommend the use of *Hypocreodendron*.

Use *Hypoxylon* Bull. 1791 (S) rather than *Nodulisporium* Preuss 1849 (A)

The genus *Hypoxylon*, typified by *H. fragiforme*, is a well-known group of stromatic ascomycetes on rotting wood. Although a number of segregated genera are now recognized such as *Annulohypoxylon*, *Biscogniauxia*, and *Nemania*, many species still remain within *Hypoxylon sensu stricto* (Hsieh *et al.* 2005). Von Arx (1982), *vide* Deighton (1985), demonstrated the relationship of the asexual morph *Nodulisporium ochraceum*, type of *Nodulisporium*, to *Hypoxylon fragiforme*, thus these generic names are synonyms. Other species of *Nodulisporium* have been placed in various genera suggesting the artificial circumscription of this genus. Given the widespread use of *Hypoxylon*, the number of names, and priority, the use of *Hypoxylon* is recommended.

Use *Microdochium* Syd. & P. Syd. 1924 (A) rather than *Monographella* Petr. later in 1924 (S)

The genus *Microdochium*, typified by *M. phragmitis*, was published a few months later than *Monographella*, typified by *M. divergens*, now regarded as a synonym of *Microdochium nivale*. Hernández-Restrepo *et al.* (2016) recognized the synonym of *Microdochium* and *Monographella* and considered *Microdochium* to be the best generic name to use for these fungi because *Microdochium* included more species and was more widely known than *Monographella*. They added six species to the 31 names in the genus *Microdochium* while *Monographella* includes only 15 names. In addition they epitypified the type of *Microdochium*, *M. phragmitis*, a species that had been shown to represent two sympatric species (Ernst *et al.* 2011). Thus, in agreement with Hernández-Restrepo *et al.* (2016), we recommend the protection of *Microdochium* over *Monographella*.

Use *Nemania* Gray 1821 (S) rather than *Geniculosporium* Chesters & Greenh. 1964 (A)

An asexual morph of *Nemania serpens*, type species of *Nemania*, was described as *Geniculosporium serpens*, type of *Geniculosporium* (Chesters & Greenhalgh 1964), thus the generic names *Nemania* and *Geniculosporium* are synonyms. *Nemania* includes over 30 names while only four names have been placed in *Geniculosporium*. In addition, *Nemania* is well-known and has priority, thus we recommend the use of *Nemania*.

Use *Rosellinia* De Not. 1844 (S) rather than *Dematophora* R. Hartig 1883 (A)

The genus *Rosellinia*, typified by *R. aquila*, is a well-known genus that has recently been monographed (Petri 2013). A number of plant pathogenic species are included in *Rosellinia* such as *R. bunodes* causing black root rot of tropical woody plants and *R. thelena* causing root collar of hardwoods. *Dematophora necatrix*, type of *Dematophora*, is considered the asexual morph of *R. necatrix*, cause of white root rot (Petri 2013), thus *Rosellinia* and *Dematophora* are synonyms. The only other species of *Dematophora*, *D. glomerata*, is considered the asexual morph of *R. glomerata*.

Given the widespread use of *Rosellinia*, the recent monographic account, its importance as a plant pathogen, and priority, we recommend the use of *Rosellinia*.

Use *Virgaria* Nees 1816 (A) rather than *Ascovirgaria* J.D. Rogers & Y.-M. Ju 2002 (S)

The monotypic genus *Ascovirgaria*, typified by *A. occulta*, was described for the sexual morph of the type species of *Virgaria*, *V. nigra*, by Rogers & Ju (2002), thus these generic names are synonyms. Nonaka *et al.* (2013) demonstrated that *A. occulta* was a synonym of *V. nigra* and that *Virgaria* was distinct from other genera in the *Xylariaceae*. Although some names have been transferred to other genera, *Virgaria* still includes 10 names and is more commonly used than *Ascovirgaria*. Given its priority and widespread use, *Virgaria* is recommended for use over *Ascovirgaria*.

Use *Xylaria* Hill ex Schrank 1789 (S) rather than *Moelleroclavus* Henn. 1902 (A), *Xylocoremium* J.D. Rogers 1984 (A), *Arthroxyllaria* Seifert & W. Gams 2002 (A) or *Geniculisynnema* Okane & Nakagiri 2007 (A)

Xylaria is a well-known genus with many species such as *Xylaria polymorpha*, commonly known as dead man's fingers, also said to cause root rot of urban trees (Proffer 1988). The asexual morphs of species of *Xylaria* have been described in four genera. The monotype species of *Moelleroclavus*, *M. penicillioides*, has a sexual morph described as *Xylaria moelleroclavus* (Rogers *et al.* 1997) and *Xylocoremium flabelliforme*, monotype species of *Xylocoremium*, is the asexual morph of *Xylaria cubensis* (Rogers 1984), now known to be *X. flabelliformis* (Ju *et al.* 2016). Both *Arthroxyllaria* and *Geniculisynnema* are monotypic genera proposed for asexual morphs not yet connected to a sexual morph but clearly within *Xylaria* based on rDNA sequences (Seifert *et al.* 2002, Okane & Nakagiri 2007). The type species of *Arthroxyllaria*, *A. elegans*, is herein placed in *Xylaria*. Although Stadler *et al.* (2013) connected *Geniculisynnema* with *Nemania*, a BLAST search with the ITS sequence of *G. termiticola* (AB274813), type species of *Geniculisynnema*, showed top matches with *Xylaria* species associated with termite nests. These four generic names are now considered synonyms of *Xylaria*. Given its widespread use, the number of species, and priority, *Xylaria* is recommended for use.

New combinations:

***Xylaria benjaminii* Seifert & W. Gams, nom. nov.**
MycBank MB816839

Replaced name: *Arthroxyllaria elegans* Seifert & W. Gams, *Czech Mycol.* **53**: 209 (2002).
Non *Xylaria elegans* Syd. & P. Syd., *Annls mycol.* **5**: 357 (1907).

***Xylaria penicilliopsis* (Henn.) Y.-M. Ju, comb. nov.**
MycBank MB816837

Basionym: *Moelleroclavus penicilliopsis* Henn., *Hedwigia* **41**: 15 (1902).

Synonym: *Xylaria moelleroclavus* J.D. Rogers *et al.*, *Mycol. Res.* **101**: 345 (1997).

Xylaria termiticola (Okane & Nakagiri) Y.-M. Ju, **comb. nov.**

MycoBank MB816838

Basionym: *Geniculisynnema termiticola* Okane & Nakagiri, *Mycoscience* **48**: 245 (2007).

Sordariomycetidae incertae sedis

Use *Brachysporium* Sacc. 1886 (A) rather than *Cryptadelphia* Réblová & Seifert 2004 (S)

Brachysporium, typified by *B. obovatum*, is a widespread and well-established asexual genus that includes 113 species and varieties. The genus *Cryptadelphia*, typified by *C. groenendalensis*, was introduced for the sexual morphs of six species of *Brachysporium* (Réblová & Seifert 2004b). The link between the sexual and asexual morphs was experimentally proven only for *B. nigrum*, which is the asexual morph of *C. groenendalensis*. The axenic cultures of *B. obovatum* and *B. polyseptatum* derived from ascospores yielded only sterile mycelium. Recently, *C. fusiformis* was described with its asexual morph regarded as *B. fusiformis*

(Markovskaja & Treigienė 2007). Based on molecular sequence data, *B. nigrum* and *B. polyseptatum* form a strongly supported monophyletic clade (Réblová & Seifert 2004b). If *B. obovatum* is congeneric with *B. nigrum*, then the generic names *Brachysporium* and *Cryptadelphia* should be treated as synonyms. Based on priority and the greater number of species, *Brachysporium* is recommended for use.

Use *Nigrospora* Zimm. 1902 (A) rather than *Khuskia* H.J. Huds. 1963 (S)

The monotype species of *Khuskia*, *K. oryzae*, was described as the sexual morph of *Nigrospora oryzae*, a well-known species of *Nigrospora*, by Hudson (1963). He also considered *N. oryzae* and *N. panici*, the type species of *Nigrospora*, to be congeneric, thus *Nigrospora* and *Khuskia* are synonyms. *Nigrospora oryzae* is reported from a variety of hosts including marine sponges (Ding et al. 2011) and as endophytes of plants (Peršoh et al. 2010) as well as causing plant diseases (Moshrefi-Zarandi et al. 2014). Given that *Nigrospora* has priority, includes fifteen names, and is more widely known than *Khuskia*, the use of *Nigrospora* is recommended.

Table 1. Names of pleomorphic genera in *Sordariomycetes* excluding *Diaporthales*, *Hypocreales* and *Magnaporthales* indicating those that are proposed for protection (including those which are asexually typified and recommended for use over sexually typified genera). For each genus the citation, type species and accepted name is given. NCF = Nomenclature Committee for Fungi.

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<i>Arthrinium</i> Kunze, in Kunze & Schmidt, <i>Mykol. Hefte</i> 1 : 9. 1817. Typus: <i>A. caricicola</i> Kunze & J.C. Schmidt 1817.	<i>Apiospora</i> Sacc. in Atti Soc. Veneto-Trent. Sci. Nat., Padova, sér. 4 4 : 85. 1875. Typus: <i>A. montagnei</i> Sacc. 1875, now regarded as <i>Arthrinium arundinis</i> (Corda) Dyko & B. Sutton 1979. <i>Pteroconium</i> Sacc. ex Grove in <i>Hedwigia</i> 55 : 146. 1914. Typus: <i>P. asteroides</i> Grove 1914, now regarded as <i>Arthrinium pterospermum</i> (Cooke & Masee) Arx 1981. <i>Scyphospora</i> L.A. Kantsch. in <i>Bolëzni Rast.</i> 17 : 87 (1928). Typus: <i>S. phyllostachydis</i> L.A. Kantsch. 1928, now regarded as <i>Arthrinium hysterinum</i> (Sacc.) P.M. Kirk 1986.	Asexual type. Protection needed by NCF.
<i>Brachysporium</i> Sacc., <i>Syll. Fung.</i> 4 : 423. 1886. Typus: <i>B. obovatum</i> (Berk.) Sacc. 1886, basionym: <i>Helminthosporium obovatum</i> Berk. 1841.	<i>Cryptadelphia</i> Réblová & Seifert in <i>Mycologia</i> 96 : 348. 2004. Typus: <i>C. groenendalensis</i> (Sacc. et al.) Réblová & Seifert 2004, basionym: <i>Zignoëlla groenendalensis</i> Sacc. et al. 1884, now regarded as <i>Brachysporium nigrum</i> (Link) S. Hughes 1958.	Asexual type. Protection needed by NCF.
<i>Calosphaeria</i> Tul. & C. Tul., <i>Select. fung. carpol.</i> 2 : 108. 1863. Typus: <i>C. princeps</i> Tul. & C. Tul. 1863.	<i>Calosphaeriophora</i> Réblová et al. in <i>Stud. Mycol.</i> 50 : 542. 2004. Typus: <i>C. pulchella</i> Réblová et al. 2004, now regarded as <i>Calosphaeria pulchella</i> (Pers.) J. Schröt. 1897.	None.
<i>Canalisporium</i> Nawawi & Kuthub. in <i>Mycotaxon</i> 34 : 477. 1989. Typus: <i>C. caribense</i> (Hol.-Jech. & Mercado) Nawawi & Kuthub. 1989, basionym: <i>Berkleasium caribense</i> Hol.-Jech. & Mercado 1984.	<i>Ascothailandia</i> Sri-indr. et al. in <i>Mycoscience</i> 51 : 414. 2010. Typus: <i>A. grenadoidea</i> Sri-indr. et al. 2010, now regarded as <i>Canalisporium grenadoideum</i> Sri-indr. et al. 2010.	Asexual type. Protection needed by NCF.

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<p>Chaetomium Kunze, in Kunze & Schmidt, Mykol. Hefte 1: 15. 1817.</p> <p>Typus: <i>C. globosum</i> Kunze 1817.</p>	<p><i>Botryotrichum</i> Sacc. & Marchal in Bull. Soc. Roy. Bot. Belgique 24: 66. 1885.</p> <p>Typus: <i>B. piluliferum</i> Sacc. & Marchal 1885, now regarded as <i>Chaetomium piluliferum</i> J. Daniels 1961. Although the synonym <i>Sepedonium albogriseum</i> Balf.-Browne 1952 provides an older name, <i>C. piluliferum</i> will be proposed for conservation.</p> <p><i>Trichocladium</i> Harz in Bull. Soc. Imp. nat. Moscou 44: 125. 1871.</p> <p>Typus: <i>T. asperum</i> Harz 1871.</p> <p><i>Humicola</i> Traaen in Nytt Mag. Natur. 52: 31. 1914.</p> <p>Typus: <i>H. fuscoatra</i> Traaen 1914.</p>	None.
<p>Chaetosphaerella E. Müll. & C. Booth in Trans. Brit. Mycol. Soc. 58: 76. 1972.</p> <p>Typus: <i>C. phaeostroma</i> (Durieu & Mont.) E. Müll. & C. Booth 1972, basionym: <i>Sphaeria phaeostroma</i> Durieu & Mont. 1846</p>	<p><i>Oedemium</i> Link, Sp. Pl., edn 4 6(1): 42. 1824.</p> <p>Typus: <i>O. atrum</i> Link 1824, now regarded as <i>Chaetosphaerella fusca</i> (Fuckel) E. Müll. & C. Booth 1972.</p> <p><i>Veramycina</i> Subram. in Kavaka 20/21: 58. 1995.</p> <p>Typus: <i>V. elegans</i> Subram. 1995, now regarded as <i>Chaetosphaerella phaeostroma</i> (Durieu & Mont.) E. Müll. & C. Booth 1972.</p>	Protection needed by NCF for <i>Chaetosphaerella</i> 1972 over <i>Oedemium</i> 1824.
<p>Chloridium Link in Mag. Gesell. Naturf. Freunde, Berlin 3: 13. 1809.</p> <p>Typus: <i>C. viride</i> Link 1805, now regarded as <i>Chloridium virescens</i> (Pers.) W. Gams & Hol.-Jech. 1976, basionym <i>Dematium virescens</i> Pers. 1794.</p>	<p><i>Gonytrichum</i> Nees & T. Nees in Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 9: 244. 1818.</p> <p>Typus: <i>G. caesium</i> Nees 1818, now regarded as <i>Chloridium caesium</i> (Nees) Réblová & Seifert 2016</p> <p><i>Melanopsammella</i> Höhn. in Ann. Mycol. 17: 121. 1920.</p> <p>Typus: <i>M. inaequalis</i> (Grove) Höhn. 1920, basionym: <i>Eriosphaeria inaequalis</i> Grove, in Berlese & Voglino 1886, now regarded as <i>Chloridium caesium</i> (Nees) Réblová & Seifert 2016.</p>	Asexual type. Protection needed by NCF.
<p>Colletotrichum Corda, Deutschl. Fl., 3(12): 41. 1837.</p> <p>Typus: <i>C. lineola</i> Corda 1832.</p>	<p><i>Glomerella</i> Spauld. & H. Schrenk in Science 17: 751. 1903.</p> <p>Typus: <i>G. cingulata</i> (Stoneman) Spauld. & H. Schrenk 1903, now regarded as <i>Colletotrichum gloeosporioides</i> (Penz.) Penz. & Sacc. 1884.</p>	Asexual type. Protection needed by NCF.
<p>Collodiscula I. Hino & Katum. in Bull. Fac. Agric. Yamaguchi Univ. 6: 55. 1955.</p> <p>Typus: <i>C. japonica</i> I. Hino & Katum. 1955.</p>	<p><i>Acanthodochium</i> Samuels <i>et al.</i> in Mycotaxon 28: 457. 1987.</p> <p>Typus: <i>A. collodisculae</i> Samuels <i>et al.</i> 1987, now regarded as <i>Collodiscula japonica</i> I. Hino & Katum. 1955.</p>	None.
<p>Coniochaeta (Sacc.) Cooke in Grevillea 16: 16. 1887, basionym: <i>Rosellinia</i> subgen. <i>Coniochaeta</i> Sacc., Syll. Fung. 1: 269. 1882.</p> <p>Typus: <i>C. ligniaria</i> (Grev.) Cooke 1887, basionym: <i>Sphaeria ligniaria</i> Grev. 1824.</p>	<p><i>Lecythophora</i> Nannf. in Svensk Skogsvårdsförening Tidskr. 3-4: 435. 1934.</p> <p>Typus: <i>L. lignicola</i> Nannf. 1934, now regarded as <i>Coniochaeta lignicola</i> (Nannf.) Z.U. Khan <i>et al.</i> 2013.</p>	None.
<p>Conioscypha Höhn. in Ann. Mycol. 2: 58. 1904.</p> <p>Typus: <i>C. lignicola</i> Höhn. 1904.</p>	<p><i>Conioscyphascus</i> Réblová & Seifert in Stud. Mycol. 50: 100. 2004.</p> <p>Typus: <i>C. varius</i> Réblová & Seifert 2004, now regarded as <i>Conioscypha varia</i> Shearer 1973.</p>	Asexual type. Protection needed by NCF.

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<p>Cordana Preuss in <i>Linnaea</i> 24: 129. 1851.</p> <p>Typus: <i>C. pauciseptata</i> Preuss 1851.</p>	<p><i>Porosphaerella</i> E. Müll. & Samuels in <i>Sydowia</i> 35: 151. 1982.</p> <p>Typus: <i>P. cordanophora</i> E. Müll. & Samuels 1982, now regarded as <i>Cordana pauciseptata</i> Preuss 1851.</p> <p><i>Pseudobotrytis</i> Krzemien. & Badura in <i>Acta Soc. Bot. Pol.</i> 23: 761. 1954.</p> <p>Typus: <i>P. fusca</i> Krzemien. & Badura 1954, now regarded as <i>Cordana terrestris</i> (Timonin) Hern.-Rest. et al. 2014.</p>	<p>Asexual type. Protection needed by NCF.</p>
<p>Corollospora Werderm. in <i>Notizbl. Bot. Gart. Berlin-Dahlem</i> 8: 248. 1922.</p> <p>Typus: <i>Corollospora maritima</i> Werderm. 1922.</p>	<p><i>Varicosporina</i> Meyers & Kohlm. in <i>Canad. J. Bot.</i> 43: 916. 1965.</p> <p>Typus: <i>V. ramulosa</i> Meyers & Kohlm. 1965, now regarded as <i>Corollospora ramulosa</i> (Meyers & Kohlm.) Abdel-Wahab 2016.</p> <p><i>Halosigmaidea</i> Nakagiri et al. in <i>Bot. Mar.</i> 52: 355. 2009.</p> <p>Typus: <i>H. luteola</i> (Nakagiri & Tubaki) Nakagiri et al. 2009, basionym: <i>Sigmaidea luteola</i> Nakagiri & Tubaki 1982, now regarded as <i>Corollospora luteola</i> Nakagiri & Tubaki 1982.</p>	<p>None.</p>
<p>Cylindrotrichum Bonord., <i>Handb. Allgem. mykol.</i>: 88 (1851).</p> <p>Typus: <i>C. oligospermum</i> (Corda) Bonord. 1851, basionym: <i>Menispora oligosperma</i> Corda 1838.</p>	<p><i>Reticulascus</i> Réblová & W. Gams in <i>Stud. Mycol.</i> 68: 180. 2011.</p> <p>Typus: <i>R. tulasneorum</i> (Réblová & W. Gams) Réblová & W. Gams 2010, basionym: <i>Chaetosphaeria tulasneorum</i> Réblová & W. Gams 1999, now regarded as <i>Cylindrotrichum oligospermum</i> (Corda) Bonord. 1851.</p>	<p>Asexual type. Protection needed by NCF.</p>
<p>Daldinia Ces. & De Not. in <i>Comment. Soc. Crittog. Ital.</i> 1(4): 197. 1863.</p> <p>Typus: <i>D. concentrica</i> (Bolton) Ces. & De Not. 1863, basionym: <i>Sphaeria concentrica</i> Bolton 1792.</p>	<p><i>Annelosporium</i> M.L. Davey in <i>Karstenia</i> 50: 3. 2010.</p> <p>Typus: <i>A. nemorosum</i> M.L. Davey 2010, now <i>Daldinia nemorosa</i> (M.L. Davey) M. Stadler et al. 2014.</p> <p><i>Versiomyces</i> Whalley & Watling in <i>Notes R. bot. Gdn Edinb.</i> 45: 401. 1989.</p> <p>Typus: <i>V. cahuchucosus</i> Whalley & Watling 1989, now regarded as <i>Daldinia cahuchucosus</i> (Whalley & Watling) M. Stadler & Læssøe 2014.</p>	<p>None.</p>
<p>Diachora Jul. Müll. in <i>Jahrb. Wiss. Bot.</i> 25: 623. 1893.</p> <p>Typus: <i>D. onobrychidis</i> (DC.) Jul. Müll. 1893, basionym: <i>Xyloma onobrychidis</i> DC. 1815.</p>	<p><i>Diachorella</i> Höhn. in <i>Hedwigia</i> 60: 192. 1918.</p> <p>Typus: <i>D. onobrychidis</i> (DC.) Höhn. 1918, basionym: <i>Xyloma onobrychidis</i> DC. 1815, now regarded as <i>Diachora onobrychidis</i> (DC) Jul. Müll. 1893.</p>	<p>None.</p>
<p>Diatrype Fr., <i>Summa veg. Scand.</i> 2: 384. 1849.</p> <p>Typus: <i>D. disciformis</i> (Hoffm.) Fr. 1849, basionym: <i>Sphaeria disciformis</i> Hoffm. 1787.</p>	<p><i>Libertella</i> Desm. in <i>Ann. Sci. Nat. (Paris) sér. 1</i> 19: 275. 1830.</p> <p>Typus: <i>L. betulina</i> Desm. 1830, now regarded as <i>Diatrype stigma</i> (Hoffm.) Fr. 1849.</p>	<p>Protection needed by NCF for <i>Diatrype</i> 1849 over <i>Libertella</i> 1830.</p>
<p>Dyrithiopsis L. Cai et al. in <i>Mycologia</i> 95: 912. 2003.</p> <p>Typus: <i>D. lakefuxianensis</i> L. Cai et al. 2003.</p>	<p><i>Monochaetiopsis</i> L. Cai et al. in <i>Mycologia</i> 95: 913. 2003.</p> <p>Typus: <i>M. lakefuxianensis</i> L. Cai et al. 2003, now regarded as <i>Dyrithiopsis lakefuxianensis</i> L. Cai et al. 2003.</p>	<p>None.</p>

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<i>Hyalotiopsis</i> Punith. in Mycol. Pap. 119 : 12. 1970. Typus: <i>H. subramanianii</i> (Agnihotr. & Luke) Punith. 1970, basionym: <i>Hyalotiella subramanianii</i> Agnihotr. & Luke 1970.	<i>Ellurema</i> Nag Raj & W.B. Kendr. in Sydowia 38 : 178. 1986. Typus: <i>E. indica</i> (Punith.) Nag Raj & W.B. Kendr. 1986, basionym: <i>Massaria indica</i> Punith. 1970, now regarded as <i>Hyalotiopsis subramanianii</i> (Agnihotr. & Luke) Punith. 1970.	Asexual type. Protection needed by NCF.
<i>Hypocreodendron</i> Henn. in Hedwigia 36 : 223. 1897. Typus: <i>H. sanguineum</i> Henn. 1897.	<i>Discoxylaria</i> J.C. Lindq. & J.E. Wright in Darwiniana 13 : 139. 1964. Typus: <i>Discoxylaria myrmecophila</i> J.C. Lindq. & J.E. Wright 1964, now regarded as <i>Hypocreodendron sanguineum</i> Henn. 1897.	Asexual type. Protection needed by NCF.
<i>Hypoxylon</i> Bull., Hist. Champ. Fr. 1 : 168. 1791. Typus: <i>H. coccineum</i> Bull. 1791, now regarded as <i>Hypoxylon fragiforme</i> (Pers.) J. Kickx f. 1835, basionym <i>Sphaeria fragiformis</i> Pers. 1794.	<i>Nodulisporium</i> Preuss in Klotzschii Herb. Viv. Mycol.: no. 1272. 1849. Typus: <i>N. ochraceum</i> Preuss 1849, now regarded as <i>Hypoxylon fragiforme</i> (Pers.) J. Kickx f. 1835.	None.
<i>Juncigena</i> Kohlm. et al. in Bot. Mar. 40 : 291. 1997. Typus: <i>J. adarca</i> Kohlm. et al. 1997.	<i>Moheitospora</i> Abdel-Wahab et al. in Mycol. Progr. 9 : 551. 2010. Typus: <i>M. fruticosae</i> Abdel-Wahab et al. 2010, now regarded as <i>Juncigena fruticosae</i> (Abdel-Wahab et al.) A.N. Mill. & Shearer 2016.	None.
<i>Knoxdaviesia</i> M.J. Wingf. et al. in Mycologia 80 : 26. 1988. Typus: <i>K. proteae</i> M.J. Wingf. et al. 1988.	<i>Gondwanamyces</i> G.J. Marais & M.J. Wingf. in Mycologia 90 : 139. 1998. Typus: <i>G. proteae</i> (M.J. Wingf. et al.) G.J. Marais & M.J. Wingf. 1998, basionym: <i>Ceratocystiopsis proteae</i> M.J. Wingf. et al. 1988, now regarded as <i>Knoxdaviesia proteae</i> M.J. Wingf. et al. 1988.	Asexual type. Protection needed by NCF.
<i>Lasiosphaeris</i> Clem., Gen. Fungi: 173. 1909. Typus: <i>L. hispida</i> (Tode) Clem. 1909.	<i>Lasiadelphia</i> Réblová & W. Gams in Fungal Divers. 46 : 82. 2011. Typus: <i>L. lasiosphaeriae</i> (W. Gams) Réblová & W. Gams 2011, now regarded as <i>Lasiosphaeris hispida</i> (Tode) Clem. 1909.	None.
<i>Mammaria</i> Ces. ex Rabenh. in Bot. Zeit. 12 : 190. 1854. Typus: <i>M. echinobotryoides</i> Ces. 1854.	<i>Pseudocercophora</i> Subram. & Sekar in J. Singapore Natl. Acad. Sci. 15 : 58. 1986. Typus: <i>P. ingoldii</i> Subram. & Sekar 1986, now regarded as <i>Mammaria echinobotryoides</i> Ces. 1854.	Asexual type. Protection needed by NCF.
<i>Melanospora</i> Corda, Icon. Fung. 1 : 24. 1837. Typus: <i>M. zamiae</i> Corda 1837.	<i>Gonatobotrys</i> Corda, Pracht-FI. Eur. Schimmelbild.: 9. 1839. Typus: <i>G. simplex</i> Corda 1839, now regarded as <i>Melanospora simplex</i> (Corda) D. Hawks. 2016.	None.
<i>Menispora</i> Pers., Mycol. Eur. 1 : 32. 1822. Typus: <i>M. glauca</i> (Link) Pers. 1822, basionym: <i>Camptosporium glaucum</i> Link in Ehrenberg 1827.	<i>Zignoëlla</i> Sacc. in Michelia 1 : 346. 1878. Typus: <i>Z. pulviscula</i> (Curr.) Sacc. 1878, now regarded as <i>Menispora caesia</i> Preuss 1851.	Asexual type. Protection needed by NCF.
<i>Menisporopsis</i> S. Hughes in Mycol. Pap. 48 : 59. 1952. Typus: <i>M. theobromae</i> S. Hughes 1952.	<i>Menisporopascus</i> Matsush. in Matsush. Mycol. Mem. 10 : 141. 2003. Typus: <i>M. kobensis</i> Matsush. 2003, now regarded as <i>Menisporopsis kobensis</i> Matsush. 2003.	Asexual type. Protection needed by NCF.
<i>Microdochium</i> Syd. & P. Syd. in Ann. Mycol. 22 (3/6): 267. Published on 15 Nov 1924. Typus: <i>M. phragmitis</i> Syd. & P. Syd. 1924.	<i>Monographella</i> Petr. in Ann. Mycol. 22 (1/2): 144. Published 20 Jun 1924. Typus: <i>M. divergens</i> (Rehm) Petr. 1924, basionym: <i>Sphaerulina divergens</i> Rehm 1913, now regarded as <i>Microdochium nivale</i> (Fr.) Samuels & I.C. Hallett 1983.	Protection needed by NCF for <i>Microdochium</i> (Jun 1924) over <i>Monographella</i> (Nov 1924).

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
Monilochaetes Halst. ex Harter in J. Agric. Res. 5 : 791. 1916. Typus: <i>M. infuscans</i> Harter 1916.	<i>Australiasca</i> Sivan. & Alcorn in Aust. Syst. Bot. 15 : 741. 2002. Typus: <i>A. queenslandica</i> Sivan. & Alcorn 2002, now regarded as <i>Monilochaetes camelliae</i> (Alcorn & Sivan.) Réblová et al. 2011.	Asexual type. Protection needed by NCF.
Nemania Gray, Nat. Arr. Brit. Pl. 1 : 516. 1821. Type: <i>N. serpens</i> (Pers.) Gray 1821.	<i>Geniculosporium</i> Chesters & Greenh. in Trans. Brit. Mycol. Soc. 47 : 400. 1964. Type: <i>G. serpens</i> Chesters & Greenh. 1964, now regarded as <i>Nemania serpens</i> (Pers.) Gray 1821.	None.
Neurospora Shear & B.O. Dodge in J. Agric. Res. 34 : 1025. 1927. Typus: <i>N. sitophila</i> Shear & B.O. Dodge 1927.	<i>Chrysonilia</i> Arx in Sydowia 34 : 16. 1981. Typus: <i>Chrysonilia sitophila</i> (Mont.) Arx 1981, basionym: <i>Penicillium sitophilum</i> Mont. 1843, now regarded as <i>Neurospora sitophila</i> Shear & B.O. Dodge 1927.	None.
Nigrospora Zimm. in Centralbl. Bakteriол. Parasitenk., 1. Abt. 8 : 220. 1902. Typus: <i>N. panici</i> Zimm. 1902.	<i>Khuskia</i> H.J. Huds. in Trans. Brit. Mycol. Soc. 46 : 358. 1963. Typus: <i>K. oryzae</i> H.J. Huds. 1963, now regarded as <i>Nigrospora oryzae</i> (Berk. & Broome) Petch 1924.	Asexual type. Protection needed by NCF.
Ophiodothella (Henn.) Höhn. in Sitzungsber. Kaiserl. Akad. Wiss., Math.-Naturwiss. Cl., Abt. 119 : 940. 1910, basionym: <i>Ophiodothis</i> subgen. <i>Ophiodothella</i> Henn. 1904. Typus: <i>O. atromaculans</i> (Henn.) Höhn. 1910, basionym: <i>Ophiodothis atromaculans</i> Henn. 1904.	<i>Acervicypeatus</i> Hanlin in Mycotaxon 37 : 380. 1990. Typus: <i>A. poriformans</i> Hanlin 1990, now regarded as <i>Ophiodothella angustissima</i> (Peck) Hanlin & M.C. González 2013.	None.
Pestalotiopsis Steyaert in Bull. Jard. Bot. État. Bruxelles 19 : 300. 1949. Typus: <i>P. guepinii</i> (Desm.) Steyaert 1949, basionym: <i>Pestalotia guepinii</i> Desm. 1840.	<i>Pestalosphaeria</i> M.E. Barr in Mycologia 67 : 188. 1975. Typus: <i>P. concentrica</i> M.E. Barr 1975, now regarded as <i>Pestalotiopsis guepinii</i> var. <i>macrotricha</i> (Kleb.) B. Sutton 1961.	Asexual type. Protection needed by NCF.
Phaeoacremonium W. Gams et al. in Mycologia 88 : 789. 1996. Typus: <i>P. parasiticum</i> (Ajello et al.) W. Gams, et al. 1996.	<i>Romellia</i> Berl., Icon. Fung. 3 : 5. 1900. Typus: <i>R. vibratilis</i> (Fr.) Berl. 1900, basionym: <i>Sphaeria vibratilis</i> Fr. 1823, now <i>Phaeoacremonium vibratilis</i> (Fr.) Gramaje et al. 2015. <i>Togninia</i> Berl., Icon. Fung. 3 : 9. 1900. Typus: <i>T. minima</i> (Tul. & C. Tul.) Berl. 1900, now regarded as <i>Phaeoacremonium minimum</i> (Tul. & C. Tul.) D. Gramaje, et al. 2015.	Protection needed by NCF for <i>Phaeoacremonium</i> 1996 over <i>Romellia</i> 1900 and <i>Togninia</i> 1900.
Plectosphaerella Kleb. in Phytopathol. Z. 1 : 43. 1929. Typus: <i>P. cucumeris</i> Kleb. 1929.	<i>Plectosporium</i> M.E. Palm et al. in Mycologia 87 : 398. 1995. Typus: <i>P. tabacinum</i> (J.F.H. Beyma) M.E. Palm et al. 1995, now regarded as <i>Plectosphaerella cucumeris</i> Kleb. 1929.	None.
Pleurostoma Tul. & C. Tul., Select. fung. Carpol. 2 : 247. 1863. Typus: <i>P. candollei</i> Tul. & C. Tul. 1863.	<i>Pleurostomophora</i> Vijaykr. et al. in Stud. Mycol. 50 : 390. 2004. Typus: <i>P. ootheca</i> Vijaykr. et al. 2004, now regarded as <i>Pleurostoma ootheca</i> (Berk. & M.A. Curtis) M.E. Barr 1985.	None.
Pleurothecium Höhn. in Ber. Deutsch. Bot. Ges. 37 : 154. 1919. Typus: <i>P. recurvatum</i> (Morgan) Höhn. 1924, basionym: <i>Acrothecium recurvatum</i> Morgan 1895.	<i>Carpoligna</i> F.A. Fernández & Huhndorf in Mycologia 91 : 253. 1999. Typus: <i>C. pleurothecii</i> F.A. Fernández & Huhndorf 1999, now regarded as <i>Pleurothecium recurvatum</i> (Morgan) Höhn. 1924.	Asexual type. Protection needed by NCF.

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<i>Polystigma</i> DC., in de Candolle & Lamarck, Fl. Franç., edn 3 6: 164. 1815. Typus: <i>P. rubrum</i> (Pers.) DC. 1815, basionym: <i>Xyloma rubrum</i> Pers. 1800.	<i>Polystigmia</i> Sacc., Syll. Fung. 3: 622. 1884. Typus: <i>P. rubra</i> (Pers.) Sacc. 1884, basionym: <i>Xyloma rubrum</i> Pers. 1800, now regarded as <i>Polystigma rubrum</i> (Pers.) DC. 1815. <i>Rhodoseptoria</i> Naumov in Bull. Soc. Mycol. Fr. 29: 278. 1913. Typus: <i>R. ussuriensis</i> Naumov 1913, now regarded as <i>Polystigma rubrum</i> (Pers.) DC. 1815..	None.
<i>Pseudothiella</i> Petr. in Hedwigia 68: 257. 1928. Typus: <i>P. hirtellae</i> (Henn.) Petr. 1928.	<i>Pseudothiopsella</i> Petr. in Hedwigia 68: 259. 1928. Typus: <i>P. hirtellae</i> Petr. 1928, now regarded as <i>Pseudothiella hirtellae</i> (Henn.) Petr. 1928.	None.
<i>Rosellinia</i> De Not. in Giorn. Bot. ital. 1: 334. 1844. Typus: <i>R. aquila</i> (Fr.) Ces. & De Not. 1844.	<i>Dematophora</i> R. Hartig in Untersuch. Forstbot. Inst. München 3: 95. 1883. Typus: <i>D. necatrix</i> R. Hartig 1883, now regarded as <i>Rosellinia necatrix</i> Berl. ex Prill. 1904	None.
<i>Seimatosporium</i> Corda, in Sturm, Deutschl. Fl., 3 Abt. 3(13): 79. 1833. Typus: <i>S. rosae</i> Corda 1833.	<i>Discostroma</i> Clem., Gen. Fung.: 50. 1909. Typus: <i>Discostroma rehmi</i> (Schnabl) Clem. 1909, basionym: <i>Curreya rehmi</i> Schnabl 1892, a synonym of <i>D. massarina</i> (Sacc.) Arx 1974, basionym: <i>Metasphaeria massarina</i> Sacc. 1884, now regarded as <i>Seimatosporium massarina</i> (Sacc.) Jaklitsch & Voglmayr 2016.	Asexual type. Protection needed by NCF.
<i>Seiridium</i> Nees, Syst. Pilze: 22. 1816. Typus: <i>S. marginatum</i> Nees 1816.	<i>Blorgiascospora</i> Shoemaker <i>et al.</i> in Canad. J. Bot. 44: 248. 1966. Typus: <i>B. marginata</i> (Fuckel) Shoemaker <i>et al.</i> 1966, basionym: <i>Massaria marginata</i> Fuckel 1873, now regarded as <i>Seiridium marginatum</i> Nees 1816.	Asexual type. Protection needed by NCF.
<i>Sphaeronaemella</i> P. Karst. in Hedwigia 23: 17. 1884. Typus: <i>S. helvella</i> (P. Karst.) P. Karst. 1884, basionym: <i>Sphaeria helvella</i> P. Karst. 1867.	<i>Gabarnaudia</i> Samson & W. Gams in Stud. Mycol. 6: 88. 1974. Typus: <i>G. betae</i> (Delacr.) Samson & W. Gams, in Samson 1974, basionym: <i>Oospora betae</i> Delacr. 1897, now regarded as <i>Sphaeronaemella betae</i> (Delacr.) Z.W. De Beer & M.J. Wingf. 2013.	None.
<i>Sporoschisma</i> Berk. & Broome, in Berkeley, Gard. Chron. 1847: 540. 1847. Typus: <i>S. mirabile</i> Berk. & Broome 1847.	<i>Melanochaeta</i> E. Müll. <i>et al.</i> in Revue Mycol. 33: 377. 1969. Typus: <i>M. hemipsila</i> (Berk. & Broome) E. Müll., <i>et al.</i> 1969, basionym: <i>Sphaeria hemipsila</i> Berk. & Broome 1873, now regarded as <i>Sporoschisma hemipsila</i> (Berk. & Broome) Zelski <i>et al.</i> 2014.	Asexual type. Protection needed by NCF.
<i>Sporoschismopsis</i> Hol.-Jech. & Hennebert in Bull. Jard. Bot. Nat. Belgique. Bruxelles 42: 385. 1972. Typus: <i>S. moravica</i> Hol.-Jech. & Hennebert 1972.	<i>Porosphaerellopsis</i> Samuels & E. Müll. in Sydowia 35: 143. 1982. Typus: <i>P. sporoschismophora</i> (Samuels & E. Müll.) E. Müll. & Samuels 1982, basionym: <i>Porosphaeria sporoschismophora</i> Samuels & E. Müll. 1978, now regarded as <i>Sporoschismopsis sporoschismophora</i> (Samuels & E. Müll.) Réblová 2014.	Asexual type. Protection needed by NCF.
<i>Stanjehughesia</i> Subram. in Proc. Indian Acad. Sci., B, Biol. Sci. 58(4): 184. 1992. Type: <i>S. hormiscioides</i> (Corda) Subram. 1992, basionym: <i>Sporidesmium hormiscioides</i> Corda 1838.	<i>Umbrinosphaeria</i> Réblová in Mycotaxon 71: 17. 1999. Type: <i>U. caesariata</i> (Clinton & Peck) Réblová 1999, basionym: <i>Sphaeria caesariata</i> Clinton & Peck, in Peck 1878, now regarded as <i>Stanjehughesia hormiscioides</i> (Corda) Subram. 1992.	Asexual type. Protection needed by NCF.

Table 1. (Continued).

Generic name recommended for use, citation and type species	Suppressed generic name(s), citation, type species and currently accepted name	Action Required
<i>Stromatographium</i> Höhn. in Denkschr. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl. 83 : 37. 1907. Typus: <i>S. stromaticum</i> (Berk.) Höhn. 1907, basionym: <i>Stilbum stromaticum</i> Berk. 1843.	<i>Fluviostroma</i> Samuels & E. Müll. in Sydowia 33 : 283. 1980. Typus: <i>F. wrightii</i> Samuels & E. Müll. 1980, now regarded as <i>Stromatographium stromaticum</i> (Berk.) Höhn. 1907.	None.
<i>Togniniella</i> Réblová et al. in Stud. Mycol. 50 : 543. 2004. Typus: <i>T. acerosa</i> Réblová et al. 2004.	<i>Phaeocrella</i> Réblová et al. in Stud. Mycol. 50 : 545. 2004. Typus: <i>P. acerosa</i> Réblová et al. 2004, now regarded as <i>Togniniella acerosa</i> Réblová et al. 2004.	None.
<i>Torpedospora</i> Meyers in Mycologia 49 : 496. 1957. Typus: <i>T. radiata</i> Meyers 1957.	<i>Glomerulispora</i> Abdel-Wahab & Nagah. in Mycol. Progr. 9 : 552. 2010. Typus: <i>G. mangrovei</i> Abdel-Wahab & Nagah. 2010, now regarded as <i>Torpedospora mangrovei</i> (Abdel-Wahab & Nagah.) E.B.G. Jones & Abdel-Wahab 2016.	None.
<i>Virgaria</i> Nees, Syst. Pilze: 54 . 1816. Typus: <i>V. nigra</i> (Link) Nees 1817, basionym: <i>Botrytis nigra</i> Link 1809.	<i>Ascovirgaria</i> J.D. Rogers & Y.M. Ju in Canad. J. Bot. 80 : 478. 2002. Typus: <i>A. occulta</i> J.D. Rogers & Y.M. Ju 2002, now regarded as <i>Virgaria nigra</i> (Link) Nees 1817.	Asexual type. Protection needed by NCF.
<i>Xylaria</i> Hill ex Schrank, Baier. Fl. 1 : 200. 1789. Typus: <i>X. hypoxylon</i> (L.) Grev. 1824, basionym: <i>Clavaria hypoxylon</i> L. 1753.	<i>Moelleroclavus</i> Henn. in Hedwigia 41 : 15. 1902. Typus: <i>M. penicilliosis</i> Henn. 1902, now regarded as <i>Xylaria penicilliosis</i> (Henn.) Y.M. Ju 2016. <i>Xylocoremium</i> J.D. Rogers in Mycologia 76 : 913. 1984. Typus: <i>X. flabelliforme</i> (Schwein.) J.D. Rogers 1984, basionym: <i>Sphaeria flabelliformis</i> Schwein., in Fries 1828, now regarded as <i>Xylaria flabelliformis</i> (Schwein.) Berk. & M.A. Curtis 1868. <i>Geniculisynnema</i> Okane & Nakagiri in Mycoscience 48 : 245. 2007. Typus: <i>G. termiticola</i> Okane & Nakagiri 2007, now regarded as <i>Xylaria termiticola</i> (Okane & Nakagiri) Y.-M. Ju 2016. <i>Arthroxyllaria</i> Seifert & W. Gams in Czech Mycol. 53 : 299. 2002. Typus: <i>A. elegans</i> Seifert & W. Gams 2002, now regarded as <i>Xylaria benjaminii</i> Seifert & W. Gams 2016.	None.

REFERENCES

- Abdel-Wahab MA, Nagahama T, Abdel-Aziz FA (2009) Two new *Corollospora* species and one new anamorph based on morphological and molecular data. *Mycoscience* **50**: 147–155.
- Abdel-Wahab MA, Pang KL, Nagahama T, Abdel-Aziz FA, Jones EBG (2010) Phylogenetic evaluation of anamorphic species of *Cirrenalia* and *Cumulospora* with the description of eight new genera and four new species. *Mycological Progress* **9**: 537–558.
- Arx JA von (1957) Die Arten der Gattung *Colletotrichum* Cda. *Phytopathologische Zeitschrift* **29**: 413–468.
- Arx JA von (1981) On *Monilia sitophila* and some families of *Ascomycetes*. *Sydowia* **34**: 13–29.
- Arx JA von (1982) The genus *Dicyma*, its synonyms and related fungi. *Proceedings, Koninklijke Nederlandse Akademie van Wetenschappen C* **85**: 21–28.
- Arzanlou M, Groenewald JZ, Gams W, Braun U, Shin HD et al. (2007) Phylogenetic and morphotaxonomic revision of *Ramichloridium* and allied genera. *Studies of Mycology* **58**: 57–93.
- Barr ME (1975) *Pestalospaeria*, a new genus in the *Amphisphaeriaceae*. *Mycologia* **67**: 187–194.
- Berbegal M, Garcia-Jimenez J, Armengol J (2014) First report of

- Calosphaeria pulchella* causing canker and branch dieback of sweet cherry trees in Spain. *Plant Disease* **98**: 1008.
- Booth C (1957) Studies of pyrenomycetes: I. Four species of *Chaetosphaeria*, two with *Catenularia* conidia. II. *Melanopsamma pomiformis* and its *Stachybotrys* conidia. *Mycological Papers* **68**: 1–27.
- Booth C (1958) The genera *Chaetosphaeria* and *Thaxteria* in Britain. *Naturalist* **1958**: 83–90.
- Braun U (2013) (2210–2232) Proposals to conserve the teleomorph-typified name *Blumeria* against the anamorph-typified name *Oidium* and twenty-two teleomorph-typified powdery mildew species names against competing anamorph-typified names (*Ascomycota*: *Erysiphaceae*). *Taxon* **62**: 1328–1331.
- Brockmann I (1976) Untersuchungen ueber die Gattung *Discostroma* Clements (*Ascomycetes*). *Sydowia* **28**: 275–338.
- Cannon PF (1991) A revision of *Phyllachora* and some similar genera on the host family *Leguminosae*. *Mycological Papers* **163**: 1–302.
- Cannon PF (1996) Systematics and diversity of the *Phyllachoraceae* associated with *Rosaceae*, with a monograph of *Polystigma*. *Mycological Research* **100**: 1409–1427.
- Cannon PF, Damm U, Johnston PR, Weir BS (2012) *Colletotrichum*—current status and future directions. *Studies in Mycology* **73**: 181–213.
- Cannon PF, Hawksworth DL (1982) A re-evaluation of *Melanospora* Corda and similar *Pyrenomycetes*, with a revision of the British species. *Botanical Journal of the Linnean Society* **84**: 115–160.
- Carlucci A, Raimondo ML, Santos J, Phillips AJL (2012) *Plectosphaerella* species associated with root and collar rots of horticultural crops in southern Italy. *Persoonia* **28**: 34–48.
- Chesters CGC, Greenhalgh GN (1964) *Geniculosporium serpens* gen. et sp. nov., the imperfect state of *Hypoxylon serpens*. *Transactions of the British Mycological Society* **47**: 393–401.
- Clements FE (1909) *The Genera of Fungi*. Minneapolis: H.W. Wilson.
- Clements FE, Shear C (1931) *The Genera of Fungi*. New York: H. W. Wilson.
- Constantinescu O, Holm K, Holm L (1995) Teleomorph-anamorph connections in ascomycetes: the anamorphs of three species of *Chaetosphaeria*. *Mycological Research* **99**: 585–592.
- Crous PW, Gams W, Wingfield MJ, van Wyk PS (1996) *Phaeoacremonium* gen. nov. associated with wilt and decline diseases of woody hosts and human infections. *Mycologia* **88**: 786–796.
- Crous PW, Groenewald JZ (2013) A phylogenetic re-evaluation of *Arthrimum*. *IMA Fungus* **4**: 133–154.
- Crous PW, Verkley GJM, Christensen M, Castaneda-Ruiz RF, Groenewald JZ (2012) How important are conidial appendages? *Persoonia* **28**: 126–137.
- Damm U, Cannon PF, Woudenberg JHC, Crous PW (2012) The *Colletotrichum acutatum* species complex. *Studies in Mycology* **73**: 37–113.
- Damm U, Fourie PH, Crous PW (2010) *Coniochaeta* (*Lecytophthora*), *Collophora* gen. nov. and *Phaeomoniella* species associated with wood necroses of *Prunus* trees. *Persoonia* **24**: 60–80.
- Damm U, Woudenberg JHC, Cannon PF, Crous PW (2009) *Colletotrichum* species with curved conidia from herbaceous hosts. *Fungal Diversity* **39**: 45–87.
- Daniel H-M, Lachance M-A, CP Kurtzman (2014) On the reclassification of species assigned to *Candida* and other ascomycetous yeast genera based on phylogenetic circumscription. *Antonie van Leeuwenhoek* **10**: 67–84.
- Daniels J (1961) *Chaetomium piluliferum* sp. nov., the perfect state of *Botryotrichum piluliferum*. *Transactions of the British Mycological Society* **44**: 79–86.
- de Beer ZW, Seifert KA, Wingfield MJ (2013) A nomenclator for ophiostomatoid genera and species in the *Ophiostomatales* and *Microascales*. *CBS Biodiversity Series* **12**: 245–322.
- Deighton FC (1985) Some species of *Nodulisporium*. *Transactions of the British Mycological Society* **85**: 391–395.
- Ding B, Yin Y, Zhang F, Li Z (2011) Recovery and phylogenetic diversity of culturable fungi associated with marine sponges *Clathrina luteoculcitella* and *Holoxea* sp. in the South China Sea. *Marine Biotechnology* **13**: 713–721.
- Ellis MB (1965) Dematiaceous Hyphomycetes. VI. *Mycological Papers* **103**: 1–46.
- Ellis MB (1971) *Dematiaceous Hyphomycetes*. Kew: Commonwealth Mycological Institute.
- Ernst M, Neubert K, Mendgen K, Wiersel SGR (2011) Niche differentiation of two sympatric species of *Microdochium* colonizing the roots of common reed, *M. phragmitis* vs. *M. bolleyi*. *BMC Microbiology* **11**: 242–255.
- Fernández FA, Huhndorf SM (2005) New species of *Chaetosphaeria*, *Melanopsammella* and *Tainosphaeria* gen. nov. from the Americas. *Fungal Diversity* **18**: 15–57.
- Fernández FA, Lutzoni FM, Huhndorf SM (1999) Teleomorph-anamorph connections: the new pyrenomycetous genus *Carpoligna* and its *Pleurothecium* anamorph. *Mycologia* **91**: 251–262.
- Fernández FA, Miller AN, Huhndorf SM, Lutzoni FM, Zoller S (2006) Systematics of the genus *Chaetosphaeria* and its allied genera: morphological and phylogenetic diversity in north temperate and neotropical taxa. *Mycologia* **98**: 121–130.
- Furlanetto C, Dianese JC (1998) Some coelomycetes from Central Brazil. *Mycological Research* **102**: 19–29.
- Gams W (2000). *Phialophora* and some similar morphologically little-differentiated anamorphs of divergent ascomycetes. *Studies in Mycology* **45**: 187–199.
- Gams W, Holubová-Jechová V (1976) *Chloridium* and some other dematiaceous hyphomycetes growing on decaying wood. *Studies in Mycology* **13**: 1–99.
- Gramaje D, Mostert L, Groenewald JZ, Crous PW (2015) *Phaeoacremonium*: from esca disease to phaeohyphomycosis. *Fungal Biology* **119**: 759–783.
- Grove WB (1937) *British Stem and Leaf-Fungi (Coelomycetes)*. Vol. 2. Cambridge: Cambridge University Press.
- Habibi A, Banihashemi Z, Mostowfzadeh-Ghalamfarsa R (2015) Phylogenetic analysis of *Polystigma* and its relationship to *Phyllachorales*. *Phytopathologia Mediterranea* **54**: 45–54.
- Hambleton S, Nickerson NL, Seifert KA (2005) *Leohumicola*, a new genus of heat-resistant hyphomycetes. *Studies in Mycology* **53**: 29–52.
- Hanlin RT (1990) *Acerviclypeatus*, a new genus for the anamorph of *Ophiodothella vaccinii*. *Mycotaxon* **37**: 379–384.
- Hanlin RT, Goh T-K, Skarshaug AJ (1992) A key to and descriptions of species assigned to *Ophiodothella* based on the literature. *Mycotaxon* **44**: 103–126.
- Hanlin RT, González M (2013) *Ophiodothella angustissima* comb. nov., a new name for *Acerviclypeatus poriformans* and *O. vaccinii*. *Mycotaxon* **123**: 327–334.
- Hausner G, Reid J (2004) The nuclear small subunit ribosomal genes of *Sphaeronaemella helvella*, *Sphaeronaemella fimicola*,

- Gabarnaudia betae*, and *Cornuvesica falcata*: phylogenetic implications. *Canadian Journal of Botany* **82**: 752–762.
- Hawksworth DL (2015) Proposals to clarify and enhance the naming of fungi under the *International Code of Nomenclature for algae, fungi, and plants*. *IMA Fungus* **6**: 199–205; *Taxon* **64**: 858–862.
- Hawksworth DL, Gibson IAS, Gams W (1976) *Phialophora parasitica* associated with disease conditions in various trees. *Transactions of the British Mycological Society* **66**: 427–431.
- Hernández-Restrepo M, Gene J, Mena-Portales J, Cano J, Hadrid H, et al. (2014) New species of *Cordana* and epitypification of the genus. *Mycologia* **106**: 723–734.
- Hernández-Restrepo M, Groenewald JZ, Crous PW (2016) Taxonomy and phylogenetic reevaluation of *Microdochium*, *Monographella* and *Idriella*. *Persoonia* **36**: 57–82.
- Hironaga M, Nakano K, Yokoyama I, Kitajima J (1989) *Phialophora repens*, an emerging agent of subcutaneous phaeohyphomycosis in humans. *Journal of Clinical Microbiology* **27**: 394–399.
- Holubová-Jechová V (1973a) Lignicolous hyphomycetes from Czechoslovakia 4. *Menispora*. *Folia Geobotanica et Phytotaxonomic* **8**: 317–336.
- Holubová-Jechová V (1973b) Lignicolous hyphomycetes from Czechoslovakia 3. *Sporoschisma*, *Sporoschismopsis* and *Catenularia*. *Folia Geobotanica et Phytotaxonomic* **8**: 209–218.
- Holubová-Jechová V, Hennebert GL (1972) *Sporoschismopsis*, a new genus of lignicolous hyphomycetes. *Bulletin du Jardin Botanique National de Belgique* **42**: 385–391.
- Hsieh H-M, Ju Y-M, Rogers JD (2005) Molecular phylogeny of *Hypoxylon* and closely related genera. *Mycologia* **97**: 844–865.
- Hudson HJ (1963) The perfect state of *Nigrospora oryzae*. *Transactions of the British Mycological Society* **46**: 355–360.
- Hughes SJ (1951) Studies on micro-fungi. XI. Some hyphomycetes which produce phialides. *Mycological Papers* **45**: 1–36.
- Hughes SJ (1958) Revisions hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Canadian Journal of Botany* **36**: 727–836.
- Hughes SL, Hennebert GL (1963) Microfungi: X. *Oedemium*, *Dimeria*, *Diplosporium*, *Gongylocladium*, and *Cladotrichum*. *Canadian Journal of Botany* **41**: 773–809.
- Jaklitsch WJ, Gardiennet, A, Voglmayr H (2016) Resolution of morphology-based taxonomic delusions: *Acrocordiella*, *Basiseptospora*, *Blogiascospora*, *Clypeosphaeria*, *Hymenopleella*, *Lepteutypa*, *Pseudapiospora*, *Requienella*, *Seiridium* and *Strickeria*. *Persoonia* **37**: 82–105.
- Jeewon R, Cai L, Liew ECY, Zhang K-Q, Hyde KD (2003) *Dyrithiopsis lakefuxianensis* gen. et sp. nov. from Fuxian Lake, Yunnan, China, and notes on the taxonomic confusion surrounding *Dyrithium*. *Mycologia* **95**: 911–920.
- Johnston PR, Seifert KA, Stone JK, Rossman AY, Maranova L (2014) Recommendations on generic names competing for use in *Leotiomycetes* (Ascomycota). *IMA Fungus* **5**: 91–120.
- Jones EBG, Suetrong S, Sakayaroj J, Bahkali AH, Abdel-Wahab MA, et al. (2015) Classification of marine Ascomycota, Basidiomycota, Blastocladiomycota and Chytridiomycota. *Fungal Diversity* **73**: 1–72.
- Jones EBG, Zuccaro A, Mitchell J, Nakagira A, Chatmala I, et al. (2009) Phylogenetic position of freshwater and marine *Sigmoidea* species: introducing a marine hyphomycete *Halosigmoidea* gen. nov. (*Halosphaeriales*). *Botanica Marina* **52**: 349–359.
- Ju Y-M, Hsieh H-M, Dominick S (2016) The *Xylaria* names proposed by C.G. Lloyd. *North American Fungi* **11**: 1–31.
- Khan Z, Gene J, Ahmad S, Cano J, Al-Swieh, N et al. (2013) *Coniochaeta polymorpha*, a new species from endotracheal aspirate of a preterm neonate, and transfer of *Lecytophora* species to *Coniochaeta*. *Antonie van Leeuwenhoek* **104**: 243–252.
- Kirk PM (1986) New or interesting microfungi XV. Miscellaneous hyphomycetes from the British Isles. *Transactions of the British Mycological Society* **86**: 409–428.
- Kohlmeyer J, Volkman-Kohlmeyer B, Eriksson OE (1997) Fungi on *Juncus roemerianus*. 9. New obligate and facultative marine Ascomycotina. *Botanica Marina* **40**: 291–300.
- Kutorga E, Rukšėnienė J, Treigienė A (2006) Microscopic fungi on *Carpinus betulus* in Lithuania 1. Teleomorphs and associated anamorphs. *Botanica Lithuanica* **12**: 233–241.
- Li Q, Wen T-C, Kang J-C, Hyde KD (2015a) A new species of *Collo-discula* (Xylariaceae) from China. *Phytotaxa* **205**: 187–196.
- Li Q, Kang J-C, Hyde KD (2015b) Two new species of the genus *Collo-discula* (Xylariaceae) from China. *Mycological Progress* **14**: article 52.
- Lindqvist JC, Wright JE (1964) *Discoxyllaria* genero nuevo la forma perfecta de *Hypocreodendron*. *Darwiniana* **13**: 138–143.
- Maharachchikumbura SSN, Guo L-D, Cai L, Chukeatirote E, Wu WP, et al. (2012) A multi-locus backbone tree for *Pestalotiopsis*, with a polyphasic characterization of 14 new species. *Fungal Diversity* **56**: 95–129.
- Maharachchikumbura SSN, Hyde KD, Groenewald JZ, Xu J, Crous PW (2014) *Pestalotiopsis* revisited. *Studies in Mycology* **79**: 121–186.
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EHC, Huang S-K, et al. (2015) Towards a natural classification and backbone tree for *Sordariomycetes*. *Fungal Diversity* **72**: 199–301.
- Markovskaja S, Treigienė A (2007) A new and a rare species of *Cryptadelphia* and their *Brachysporium* anamorphs. *Nova Hedwigia* **84**: 495–501.
- McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, et al. (2006) *International Code of Botanical Nomenclature (Vienna Code)*. [Regnum Vegetabile no. 146.] Ruggell: A. R. G. Gantner Verlag.
- McNeill J, Barrie FF, Buck WR, Demoulin V, Greuter W, et al. (eds) (2012) *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code)*. [Regnum vegetabile no. 154.] Königstein: Koeltz Scientific Books.
- Meyer WM, Dooley JR, Kwon-Chung KJ (1975) Mycotic granuloma caused by *Phialophora repens*. *American Journal of Clinical Pathology* **64**: 549–555.
- Mhmoud NA, Abdalla AS, Fahal AH, de Hoog GS, Gerrits van den Ende AH et al. (2012) *Pleurostomophora ochracea*, a novel agent of human eumycetoma with yellow grains. *Journal of Clinical Microbiology* **50**: 2987–2994.
- Miller AN, Huhndorf SM (2004). A natural classification of *Lasiosphaeria* based on nuclear LSU rDNA sequences. *Mycological Research* **108**: 26–34.
- Moshrefi-Zarandi D, Aminae MM, Sharzei A, Rezaee S (2014) First report of rosemary leaf spot caused by *Nigrospora oryzae* in Iran. *New Disease Reports* **30**: 27.
- Mostert L, Crous PW, Groenewald JZ, Gams W, Summerbell RC (2003) *Togninia* (*Calosphaeriales*) is confirmed as teleomorph of *Phaeoacremonium* by means of morphology, sexual compatibility and DNA phylogeny. *Mycologia* **95**: 646–659.

- Mostert L, Groenewald JZ, Summerbell RC, Gams W, Crous PW (2006) Taxonomy and pathology of *Togninia* (*Diaporthales*) and its *Phaeoacremonium* anamorphs. *Studies in Mycology* **54**: 1–115.
- Mugambi G, Huhndorf SM (2008) A new species of *Melanochaeta* from Kenya. *Sydowia* **60**: 261–266.
- Mugambi GK, Huhndorf SM (2010) Multigene phylogeny of the *Coronophorales*: morphology and new species in the order. *Mycologia* **102**: 185–210.
- Müller E (1986) On the genus *Diachora* J. Müller. *Transactions of the Botanical Society of Edinburgh* **45** (Supplement 1): 69–75.
- Müller E, von Arx JA (1962) Die Gattungen der didymosporenen Pyrenomyceten. *Beiträge zur Kryptogamenflora der Schweiz* **11**(2): 1–922.
- Müller E, Samuels GJ (1982) Anamorphs of pyrenomycetous ascomycetes I. *Rhamphoria* Niessl and *Trichosphaerella* Bommer *et al.* *Sydowia* **35**: 143–149
- Müller E, Samuels G (1982) Anamorphs of pyrenomycetous ascomycetes. II. *Porosphaerella* gen. nov. and its *Cordana* anamorph. *Sydowia* **35**: 150–154.
- Nag Raj TR (1975) Genera coelomycetum. XI. *Hyalotia*, *Hyalotiella*, and *Hyalotiopsis*. *Canadian Journal of Botany* **53**: 1615–1624.
- Nag Raj TR (1993) *Coelomycetous Anamorphs with Appendage-bearing Conidia*. Waterloo: Mycologue Publications.
- Nag Raj TR, Kendrick B (1986) *Ellurema* gen. nov., with notes on *Lepteutypa cisticola* and *Seiridium canariense*. *Sydowia* **38**: 178–193.
- Nakagiri A, Tokura R (1982) A new marine ascomycete and its anamorph from Japan. *Transactions of the Mycological Society of Japan* **23**: 101–110.
- Nawawi A, Kuthubutheen AJ (1989) *Canalisporium*, a new genus of lignicolous hyphomycetes from Malaysia. *Mycotaxon* **34**: 475–487.
- Nonaka K, Ishit T, Shiomi K, Omura S, Masume R (2013) *Virgaria boninensis*, a new hyphomycete (*Xylariaceae*) from soils in the Bonin Islands, Japan. *Mycoscience* **54**: 394–399.
- Nonaka K, Miyazaki H, Iwatsuki M, Shiomi K, Tomoda H *et al.* (2012) *Staphylotrichum boninense*, a new hyphomycete (*Chaetomiaceae*) from soils in the Bonin Islands, Japan. *Mycoscience* **53**: 312–318.
- Norphanphoun C, Maharachchikumbura SSN, Daranagama A, Bulgakov TS, Bhat DJ, *et al.* (2015) Towards a backbone tree for *Seimatosporium*, with *S. physocarpi* sp. nov. *Mycosphere* **6**: 385–400.
- Okane I, Nakagiri A (2007) Taxonomy of an anamorphic xylariaceous fungus from a termite nest found together with *Xylaria angulosa*. *Mycoscience* **48**: 240–249.
- Palm ME, Gams W, Nirenberg HI (1995) *Plectosporium*, a new genus for *Fusarium tabacinum*, the anamorph of *Plectosphaerella cucumerina*. *Mycologia* **87**: 397–406.
- Perdomo H, Garcia D, Gene J, Sutton DA, Summerbell R *et al.* (2013) *Phialemoniopsis*, a new genus of *Sordariomycetes*, and new species of *Phialemonium* and *Lecythophora*. *Mycologia* **105**: 398–421.
- Peršoh D, Melcher M, Flessa F, Rambold G (2010) First fungal community analyses of endophytic ascomycetes associated with *Viscum album* ssp. *austriacum* and its host *Pinus sylvestris*. *Fungal Biology* **114**: 585–596.
- Petrini L (2013) *Rosellinia* – a world monograph. *Bibliotheca Mycologica* **205**: 1–410.
- Proffer TJ (1988) *Xylaria* root rot of urban trees caused by *Xylaria polymorpha*. *Plant Disease* **72**: 79.
- Punithalingam E (1970) Studies on *Sphaeropsidales* in culture. *Mycological Papers* **119**: 1–24.
- Quandt CA, Kepler RM, Gams W, Araújo JPM, Ban S, *et al.* (2014) Phylogenetic-based nomenclatural proposals for *Ophiocordycipitaceae* (*Hypocreales*) with new combinations in *Tolypocladium*. *IMA Fungus* **5**: 121–134.
- Raja HA, Fournier J, Shearer CA, Miller AN (2012) Freshwater ascomycetes: *Coniochaeta gigantospora* sp. nov. based on morphological and molecular data. *Mycoscience* **53**: 373–380.
- Rambelli A, Onofri S (1987) New species of *Kylindria* and *Xenokylindria* and notes on *Cylindrotrichum* (*Hyphomycetes*). *Transactions of the British Mycological Society* **88**: 393–397.
- Ranghoo VM, Tsui CKM, Hyde KD (2001) *Brunneospora aquatica* gen. et sp. nov., *Aqualignicola hyalina* gen. et sp. nov., *Jobellisia viridifusca* sp. nov. and *Porosphaerellopsis bipolaris* sp. nov. (*Ascomycetes*) from submerged wood in freshwater habitats. *Mycological Research* **105**: 625–633.
- Réblová M (1999) Studies in *Chaetosphaeria sensu lato* I. The genera *Chaetosphaerella* and *Tengiomyces* gen. nov. of the *Helminthosphaeriaceae*. *Mycotaxon* **70**: 387–420.
- Réblová M (2000) The genus *Chaetosphaeria* and its anamorphs. *Studies in Mycology* **45**: 149–168.
- Réblová M (2011) New insights into the systematics and phylogeny of the genus *Jattaea* and similar fungi of the *Calosphaeriales*. *Fungal Diversity* **49**: 167–198.
- Réblová M (2014) *Sporoschismopsis angustata* sp. nov., a new holomorph species in the *Reticulascaceae* (*Glomerellales*), and a reappraisal of *Sporoschismopsis*. *Mycological Progress* **13**: 671–681.
- Réblová M, Barr ME, Samuels GJ (1999) *Chaetosphaeriaceae*, a new family for *Chaetosphaeria* and its relatives. *Sydowia* **51**: 49–70.
- Réblová M, Gams W (1999) Teleomorph-anamorph connections in ascomycetes. 1. *Cylindrotrichum* and *Cacumisporium* anamorphs of *Chaetosphaeria*. *Czech Mycology* **51**: 1–40.
- Réblová M, Gams W, Seifert KA (2011a) *Monilochaetes* and allied genera of the *Glomerellales*, and a reconsideration of families in the *Microascales*. *Studies in Mycology* **68**: 163–191.
- Réblová M, Gams W, Štěpánek V (2011b) The new hyphomycete genera *Brachyalaria* and *Infundichalara*, the similar *Exochalara* and species of *Phialophora* sect. *Catenulatae* (*Leotiomyces*). *Fungal Diversity* **46**: 67–86.
- Réblová M, Jaklitsch WM, Réblová K, Štěpánek V (2015) Phylogenetic reconstruction of the *Calosphaeriales* and *Togniniales* using five genes and predicted RNA secondary structures of ITS, and *Flabellascus tenuirostris* gen. et sp. nov. *PLoS One* **10**(12): e0144616.
- Réblová M, Mostert L, Gams W, Crous PW (2004) New genera in the *Calosphaeriales*: *Togniniella* and its anamorph *Phaeocrella*, and *Calosphaeriophora* as anamorph of *Calosphaeria*. *Studies in Mycology* **50**: 533–550.
- Réblová M, Mostert L (2007) *Romellia* is congeneric with *Togninia* and description of *Conidiotheca* gen. nov. for one species of this genus with polysporous asci. *Mycological Research* **111**: 299–307.
- Réblová M, Seifert KA (2004a) *Conioscyphascus*, a new ascomycetous genus for holomorphs with *Conioscypha* anamorphs. *Studies in Mycology* **50**: 95–108.
- Réblová M, Seifert KA (2004b) *Cryptadelphia* (*Trichosphaeriales*), a new genus for holomorphs with *Brachysporium* anamorphs, and clarification of the taxonomic status of *Wallrothiella*. *Mycologia* **96**: 343–367.

- Réblová M, Seifert KA (2007) A new fungal genus, *Teracosphaeria*, with a phialophora-like anamorph (*Sordariomycetes*, *Ascomycota*). *Mycological Research* **111**: 287–298.
- Réblová M, Seifert KA, Fournier J, Štěpánek V (2016) Newly recognised lineages of perithecial ascomycetes: the new orders *Conioscyphales* and *Pleurotheciales*. *Persoonia* **37**: 57–81.
- Réblová M, Winka K (2000) Phylogeny of *Chaetosphaeria* and its anamorphs based on morphological and molecular data. *Mycologia* **92**: 939–954.
- Rodrigues de Cruz MC, Marques MFO, Guzmão LFP (2014) Conidial fungi from the semi-arid catinga biome of Brazil. The genus *Menisporopsis*. *Acta Botanica Brasiliica* **28**: 339–345.
- Rogers JD (1984) *Xylaria cubensis* and its anamorph *Xylocoremium flabelliforme*, *Xylaria allantoidea*, and *Xylaria poitei* in continental United States. *Mycologia* **76**: 912–923.
- Rogers JD, Ju Y-M (2002) *Ascovirgaria occulta* gen. et sp. nov., *Jumillera hawaiiensis* sp. nov., and *Lopadostoma hawaiianum* sp. nov. from Hawaii. *Canadian Journal of Botany* **80**: 478–481.
- Rogers JD, Ju Y-M, Hemmes DE (1997) *Xylaria moelleroclavus* sp. nov. and its *Moelleroclavus* anamorphic state. *Mycological Research* **101**: 345–348.
- Rogers JD, Ju Y-M, San Martin Gonzalez F (1995) *Discoxyllaria myrmecophila* and its *Hypocreodendron* anamorph. *Mycologia* **87**: 41–45.
- Rojas EI, Rehner SA, Samuels GJ, Van Bael SA, Herre EA et al. (2010) *Colletotrichum gloeosporioides* s.l. associated with *Theobroma cacao* and other plants in Panama: multilocus phylogenies distinguish host-associated pathogens from asymptomatic endophytes. *Mycologia* **102**: 1318–1338.
- Rolshausen PE, Mahoney NE, Molyneux RJ, Gubler WD (2006) A reassessment of the species concept in *Eutypa lata*, the causal agent of Eutypa dieback of grapevine. *Phytopathology* **96**: 367–377.
- Rossmann AY, Seifert KA, Samuels GJ, Minnis AM, Schroers H-J, et al. (2013) Genera in *Bionectriaceae*, *Hypocreaceae*, and *Nectriaceae* (*Hypocreales*) proposed for acceptance or rejection. *IMA Fungus* **4**: 41–51.
- Rossmann AY, Adams GC, Cannon PF, Castlebury LA, Crous PW et al. (2015a) Recommendations of generic names in *Diaporthales* competing for protection or use. *IMA Fungus* **6**: 145–154.
- Rossmann AY, Crous PW, Hyde KD, Hawksworth DL, Aptroot A, et al. (2015b) Recommended names of pleomorphic genera in *Dothideomycetes*. *IMA Fungus* **6**: 507–523.
- Samson RA (1974) Paecilomyces and some allied hyphomycetes. *Studies in Mycology* **6**: 1–119.
- Samson RA, Visagie CM, Houbbraken J, Hong S-B, Hubka V, et al. (2014) Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology* **78**: 141–173.
- Samuels GJ, Müller E (1978) Life-history studies of Brazilian ascomycetes. 1. Two new genera of the *Sphaeriaceae* having, respectively, *Sporoschisma*-like and *Codinaea* anamorphs. *Sydowia* **31**: 126–136.
- Samuels GJ, Müller E (1980) Life history studies of Brazilian ascomycetes 8. *Thamnomycetes chordalis* (anam.: *Nodulisporium*) and *Camillea bacillum* (anam.: *Geniculosporium*) with notes on taxonomy of the *Xylariaceae*. *Sydowia* **33**: 274–281.
- Samuels GJ, McKenzie EHC, Buchanan DE (1981) Ascomycetes of New Zealand 3. Two new species of *Apiospora* and their *Arthrimum* anamorphs on bamboo. *New Zealand Journal of Botany* **19**: 137–149.
- Samuels GJ, Rogers JD, Nagasawa E (1987) Studies in the *Amphisphaeriaceae* (*sensu lato*) 11. *Collodiscula japonica* and its anamorph, *Acanthodochium collodisculae*. *Mycotaxon* **28**: 453–459.
- Schoch CL, Sung GH, Volkmann-Kohlmeyer B, Kohlmeyer J, Spatafora JW (2007) Marine fungal lineages in the *Hypocreomycetidae*. *Mycological Research* **111**: 154–162.
- Schol-Schwarz MB (1970). Revision of the genus *Phialophora*. *Persoonia* **6**: 59–94.
- Seifert KA (1987) *Stromatographium* and *Acrostroma* gen. nov.: two tropical hyphomycete genera with distinctive synnema anatomies. *Canadian Journal of Botany* **65**: 2196–2201.
- Seifert KA, Gams W, Louis-Seize GW (2002) *Arthroxyllaria elegans*, a new coprophilous anamorphic fungus allied with the *Xylariaceae*, with notes on the genus *Bisporostilbella*. *Czech Mycology* **53**: 297–307.
- Senanayake IC, Maharachchikumbura SSN, Hyde KD, Bhat JD, Jones EBG, et al. (2015) Towards unraveling relationships in *Xylariomycetidae* (*Sordariomycetes*). *Fungal Diversity* **73**: 73–144.
- Shenoy BD, Jeewon R, Wu WP, Bhat DJ, Hyde KD (2006) Ribosomal and RPB2 DNA sequence analyses suggest that *Sporidesmium* and morphologically similar genera are polyphyletic. *Mycological Research* **110**: 916–928.
- Shoemaker RA, Müller E, Morgan-Jones G (1966) Fuckel's *Massaria marginata* and *Seiridium marginatum* Nees ex Steudel. *Canadian Journal of Botany* **44**: 247–254.
- Shoemaker RA (1964) *Seimatosporium* (= *Cryptostictis*) parasites of *Rosa*, *Vitis*, and *Cornus*. *Canadian Journal of Botany* **42**: 411–421.
- Sivanesan A (1983) Studies on ascomycetes. *Transactions of the British Mycological Society* **81**: 313–332.
- Sivichai S, Hywel-Jones NL, Somrithipol S (2000) Lignicolous freshwater *Ascomycota* from Thailand: *Melanochaeta* and *Sporoschisma* anamorphs. *Mycological Research* **104**: 478–485.
- Sri-indrasudthi V, Boonyuen N, Suetrong S, Chuaseeharonnachai C, Sivichai S, et al. (2010) Wood-inhabiting freshwater fungi from Thailand: *Ascothailandia grenadoidia* gen. et sp. nov., *Canalisporium grenadoidia* sp. nov. with a key to *Canalisporium* species (*Sordariomycetes*, *Ascomycota*). *Mycoscience* **51**: 411–420.
- Stadler M, Kuhnert E, Peršoh D, Fournier J (2013) The *Xylariaceae* as model example for a unified nomenclature following the “One fungus-one name” (1F1N) concept. *Mycology* **4**: 5–21.
- Stadler M, Læssøe T, Fournier J, Decock C, Schmiechek B, et al. (2014). A polyphasic taxonomy of *Daldinia* (*Xylariaceae*). *Studies in Mycology* **77**: 1–143.
- Subramanian CV, Sekar G (1986) *Pseudocercophora ingoldii* gen. et sp. nov. and its *Mammaria* anamorph. *Journal of the Singapore National Academy of Sciences* **15**: 58–60.
- Sutton BC (1980) *The Coelomycetes; Fungi Imperfecti with pycnidia, acervuli and stromata*. Kew: Commonwealth Mycological Institute.
- Tanaka K, Endo M, Hirayama K, Okane I, Hosoya T, et al. (2011) Phylogeny of *Discosia* and *Seimatosporium* and introduction of *Adisciso* and *Immersidiscosia* genera nova. *Persoonia* **26**: 85–98.
- Tang AM, Jeewon R, Hyde KD (2007) Phylogenetic utility of protein (RPB2, beta-tubulin) and ribosomal (LSU, SSU) gene sequences in the systematics of *Sordariomycetes*, *Ascomycota*, *Fungi*. *Antonie van Leeuwenhoek* **91**: 327–349.
- Trouillas FP, Lorber JD, Peduto F, Grant J, Coates WW, et al. (2010a)

- First report of *Calosphaeria pulchella* associated with branch dieback of sweet cherry trees in California. *Plant Disease* **94**: 1167.
- Trouillas FP, Urbez-Torres JR, Gubler WD (2010b). Diversity of diatrypaceous fungi associated with grapevine canker diseases in California. *Mycologia* **102**: 319–336.
- Untereiner WA, Debois V, Naveau FA (2001) Molecular systematics of the ascomycete genus *Farrowia* (*Chaetomiaceae*). *Canadian Journal of Botany* **79**: 321–333.
- Vakili NG (1989) *Gonatobotrys simplex* and its teleomorph, *Melanospora damnosa*. *Mycological Research* **93**: 67–74.
- Vijaykrishna D, Mostert L, Jeewon R, Gams W, Hyde KD, *et al.* (2004) *Pleurostomophora*, an anamorph of *Pleurostoma* (*Calosphaeriales*), a new anamorph genus morphologically similar to *Phialophora*. *Studies in Mycology* **50**: 387–395.
- Visagie CM, Houbraken J, Frisvad JC, Hong S-B, Klaassen CHW, *et al.* (2014) Identification and nomenclature of the genus *Penicillium*. *Studies in Mycology* **78**: 343–371.
- Weber E, Gorke C, Begerow D (2002) The *Lecythophora-Coniochaeta* complex II. Molecular studies based on sequences of the large subunit of ribosomal DNA. *Nova Hedwigia* **74**: 187–200.
- Weir BS, Johnston PR, Damm U (2012) The *Colletotrichum gloeosporioides* species complex. *Studies in Mycology* **73**: 115–180.
- Wijayawardene DNN, McKenzie EHC, Hyde KD (2012) Towards incorporating anamorphic fungi in a natural classification–checklist and notes for 2011. *Mycosphere* **3**: 157–228.
- Zelski S, Do C, Balto JA, Raja HA, Miller AN, *et al.* (2014a) Some dematiaceous freshwater microfungi from Perú and their previously unknown phylogenetic relationships based on 28S nrDNA. *IMA Fungus* **5**: 425–438.
- Zelski SE, Raja HA, Miller AN, Shearer CA (2014b) *Conioscypha peruviana* sp. nov., its phylogenetic placement based on 28S rRNA gene, and a report of *Conioscypha gracilis* comb. nov. from Peru. *Mycoscience* **56**: 319–325.
- Zhang N, Luo J, Rossman A, Aoki T, Chuma I, *et al.* (2016) Generic names in *Magnaporthales*. *IMA Fungus* **7**: 155–159.