

***Togninia* (Calosphaerales) is confirmed as teleomorph of *Phaeoacremonium* by means of morphology, sexual compatibility and DNA phylogeny**

Lizel Mostert

Department of Plant Pathology, University of Stellenbosch, P. Bag X1, Matieland 7602, South Africa

Pedro W. Crous¹

Centraalbureau voor Schimmelcultures, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands

J. Z. (Ewald) Groenewald

Department of Plant Pathology, University of Stellenbosch, P. Bag X1, Matieland 7602, South Africa

Walter Gams

Richard C. Summerbell

Centraalbureau voor Schimmelcultures, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands

Abstract: Petri disease, or black goo, is a serious disease of vines in most areas where grapevines are cultivated. The predominant associated fungus is *Phaeo-*monielliella chlamydospora** (Chaetothyriales). Several species of *Phaeoacremonium* (*Pm.*) also are associated, of which *Pm. aleophilum* is the most common. Although no teleomorph is known for *Phaeoacremonium*, the genus *Togninia* previously has been linked to phaeoacremonium-like anamorphs. To investigate the possible anamorph-teleomorph connection of *Phaeoacremonium* to *Togninia*, anamorphs of *Togninia minima*, *T. fraxinopennsylvanica* and *T. novae-zealandiae* morphologically were compared with *Pm. aleophilum* and some representative cultures were mated in all combinations. Although no interspecies mating proved fertile, matings between isolates of *Pm. aleophilum* produced a *Togninia* teleomorph within 3–4 weeks. Certain field isolates of *Pm. aleophilum* commonly produced the teleomorph, demonstrating that both mating types can occur in the same vine and thus also explaining the genetic diversity observed for this fungus in some vineyards. To elucidate the phylogenetic relationships among these taxa, isolates were subjected to sequence analysis of the nuclear ribosomal internal transcribed spacers (ITS1, ITS2) and the 5.8S rRNA gene, as well as portions of the translation elongation factor 1 alpha (EF-

1 α) gene. The generic placement of teleomorphs within *Togninia* (Calosphaerales) further was confirmed via phylogenetic analyses of 18S small subunit (SSU) DNA. From these sequences, morphological and mating data, we conclude that *T. minima* is the teleomorph of *Pm. aleophilum*, and that it has a diallelic heterothallic mating system. An epitype and mating type tester strains also are designated for *T. minima*.

Key words: Calosphaerales, EF-1 α , ITS, 18S SSU DNA, sexual compatibility, systematics

INTRODUCTION

Petri disease is a well-known disease of grapevines worldwide (Mugnai et al 1999). Affected grapevines exhibit a slow dieback as well as stunted growth. The predominant associated fungus is *Phaeo-*monielliella chlamydospora** W. Gams, Crous, M.J. Wingf. & L. Mugnai (Chaetothyriales, Herpotrichiellaceae). Whether this fungus is the sole or only a contributing causal agent of the disease is uncertain. Several species of *Phaeoacremonium* (*Pm.*) (Diaporthales, Magnaporthaceae) also commonly grow from vines affected by Petri disease, including *Pm. aleophilum* W. Gams, Crous, M.J. Wingf. & L. Mugnai, *Pm. angustius* W. Gams, Crous & M.J. Wingf., *Pm. inflatipes* W. Gams, Crous & M.J. Wingf., *Pm. mortoniae* Crous & W. Gams, *Pm. parasiticum* (Ajello, Georg & C.J.K. Wang) W. Gams, Crous & M.J. Wingf., *Pm. rubrigenum* W. Gams, Crous & M.J. Wingf., and *Pm. viticola* Dupont. Of the *Phaeoacremonium* species associated with Petri disease, *Pm. aleophilum* consistently has been the most frequently isolated (Scheck et al 1998, Ari 2000, Gatica et al 2001). In various studies, *Pm. aleophilum* has been found to be associated with brown-streaking symptoms in Petri-diseased grapevines (Mugnai et al 1999, Ari 2000) and sectorial brown necrosis (Gatica et al 2001).

The genetic variation within populations of *Phaeo-*monielliella chlamydospora** and *Pm. aleophilum* has been studied by various workers (Péros et al 2000, Tegli et al 2000a, b). Using RAPDs (Random Amplified Polymorphic DNA) and RAMS (Random Amplified Micro- or Mini-Satellites), Tegli et al (2000b) showed that considerable variation existed among isolates of *Pm. aleophilum* collected from the same field, sug-

gesting that sexual reproduction might occur (Tegli 2000). Considerable genetic variation suggestive of ongoing recombination also was found in Universally Primed-PCR studies done with *Pm. aleophilum* isolates from Australia (Cottral et al 2001). Rooney et al (2002) subsequently reported inducing teleomorphs for *Pm. inflatipes* and *Pm. aleophilum* under laboratory conditions.

In a study of the genus *Togninia* Berl. (Calosphaeriales), Hausner et al (1992) treated *Togninia minima* (Tul. & C. Tul.) Berl. (lectotype of *Togninia*) and at the same time they described two new species, *T. fraxinopennsylvanica* (Hinds) Hausner, Eyjólfsdóttir & J. Reid and *T. novae-zealandiae* Hausner, Eyjólfsdóttir & J. Reid. Although no cultures of *T. minima* were available, the two newly described species were cultured and were shown to produce anamorphs that were intermediate between *Acronium* Link : Fr. and *Phialophora* Medlar. Several anamorph genera in recent years have been described with this approximate appearance (Gams 2000). Of these, the genus *Phaeoacremonium* W. Gams et al closely fits the description of the *Togninia* anamorphs illustrated by Hausner et al (1992).

The first aim of the current study was to investigate the possible link between *Phaeoacremonium* and *Togninia*. *Phaeoacremonium aleophilum*, a species similar in appearance to the anamorphs illustrated by Hausner et al (1992), was chosen for morphological comparison with the anamorphs of *T. minima*, *T. fraxinopennsylvanica* and *T. novae-zealandiae*. A further aim was to determine if the formation of a teleomorph could be elicited in *Pm. aleophilum*. This was done by mating a selection of vine isolates in culture. A final aim was to elucidate the genetic diversity within and among these species and to clarify the higher-order phylogenetic placement of *Togninia*. To address this, a phylogenetic analysis was conducted using sequences of the nuclear ribosomal DNA region encompassing the internal transcribed spacers (ITS1, 5.8S and ITS2), as well as translation elongation factor 1 alpha (EF-1 α), and 18S ribosomal small subunit (SSU).

MATERIALS AND METHODS

Morphology.—Field isolations were made from rooted nursery plants and older, diseased grapevines, from which single-conidial isolates were obtained (TABLE I). Anamorph morphology was studied on 2% malt-extract agar (MEA; Biolab, Midrand, South Africa), while perithecia were induced on twice-autoclaved pieces of grapevine cane placed on 2% water agar (Biolab) (GWA). Cultures were incubated at 22 C under a 12 h fluorescent white light/dark regime. For microscopy, material was mounted in lactic acid. Thirty

measurements were taken of each type of morphological structure, and averages and 95% confidence intervals were determined for spore dimensions. Measurements are given with minimum and maximum ranges in parentheses.

Vertical sections (10 μ m) of fruiting bodies were cut with a Leica CM1100 freezing microtome. Colony colors were determined according to Rayner (1970). Cultures are maintained in the collection of the Department of Plant Pathology at the University of Stellenbosch, and representative strains have been deposited at the Centraalbureau voor Schimmelcultures (CBS, Utrecht, the Netherlands). Reference strains of *T. fraxinopennsylvanica* (CBS 110212, ex-type), *T. novae-zealandiae* (UAMH 9589, UAMH 9590, ex-type) and *T. minima* (CBS 213.31, ex-type of *Longoa paniculata* Curzi) also were studied.

Matings.—Twenty-one *Pm. aleophilum* isolates were grown on MEA plates for 2 wk, using 10 plates per isolate. Conidia were dislodged from the agar surface by means of a glass rod, and suspensions were prepared in 5 mL sterile distilled water. Two aliquots of 100 μ L each, representing two different isolates, were pipetted onto the canes of GWA plates. Isolates were mated in all possible combinations. Controls consisted of a 200 μ L aliquot of one isolate only. Plates were incubated at 22 C under continuous white light. Successful crosses were noted 3–4 wk after mating. For a mating to be considered successful, perithecia had to produce large quantities of ascospores that germinated readily in culture. One such mating was chosen (LM 54 \times LM 463), and 20 single ascospore isolates obtained (LM 227–LM 240, LM 243–LM 247, LM 249). Further crosses were made with these ascospore isolates using the procedure described above. Two strains found to be of opposite mating type arbitrarily were designated as MAT1-1 (LM 463) and MAT1-2 (LM 54). Inter-species matings were done to investigate the biological species boundaries of *Pm. aleophilum*, *T. minima*, *T. novae-zealandiae* and *T. fraxinopennsylvanica*.

DNA isolation and amplification.—Twenty-seven *Pm. aleophilum* isolates were selected for sequence comparisons (TABLE I). Sequences of the ITS, EF-1 α and SSU of *T. fraxinopennsylvanica* (CBS 110212), *T. novae-zealandiae* (UAMH 9589 and UAMH 9590), *T. minima* (CBS 213.31) and an unknown *Phaeoacremonium* sp. (STE-U 3394) also were included. Genomic DNA was extracted using the isolation protocol of Lee and Taylor (1990). In studies intended to determine the degree of genetic diversity within *Pm. aleophilum*, the 5.8S nuclear ribosomal RNA gene and the flanking internal transcribed spacers (ITS1 and ITS2) were amplified with primers ITS1 and ITS4 (White et al 1990) and translation elongation factor 1 alpha (EF-1 α) was amplified with primers EF1-728F and EF1-986R (Carbone and Kohn 1999). These PCR amplification cycles run on a GeneAmp PCR System 2700 (Perkin-Elmer, Norwalk, Connecticut) were for both regions: 96 C for 5 min, followed by 36 cycles of (1) denaturation (94 C for 30 s), (2) annealing (50 C for 30 s) and (3) elongation (72 C for 90 s), and a final 7 min extension step at 72 C.

In studies intended to determine the higher order phylogeny of *Togninia*, primers NS1 and NS4 (White et al 1990) were used to amplify the 5' end of the 18S ribosomal

TABLE I. *Phaeoacremonium* and *Togninia* isolates studied

Species	Culture No.	GenBank No.			Host and location
		ITS	EF	SSU	
<i>Togninia minima</i> (Anamorphi: <i>Pm. aleophilum</i>)	LM44 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM45 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM46 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM47 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM48 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM49 = CBS 110701 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM50 = CBS 110831 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM51 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM53 = CBS 110702 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM54 = CBS 110703 ^{a,c}				<i>Vitis vinifera</i> , Paarl, South Africa
	LM56 = CBS 110705 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM58 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM61 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM65 = CBS 110711 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM76 = CBS 110827 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM77 ^c				<i>Vitis vinifera</i> , Paarl, South Africa
	LM78 ^b				<i>Vitis vinifera</i> , Paarl, South Africa
	LM467 ^c				<i>Vitis vinifera</i> , Stellenbosch, South Africa
	LM468 = CBS 110753 ^b				<i>Vitis vinifera</i> , Stellenbosch, South Africa
	LM458 = CBS 111014 ^c				<i>Vitis vinifera</i> , Stellenbosch, South Africa
	LM463 = CBS 111015 ^{a,b}				<i>Vitis vinifera</i> , Stellenbosch, South Africa
	LM4 = CBS 110707	AY179930	AY179896		<i>Vitis vinifera</i> , Wellington, South Africa
	LM5 = CBS 110830	AY179935	AY179901		<i>Vitis vinifera</i> , Wellington, South Africa
	LM12 = CBS 110828	AY179929	AY179895		<i>Vitis vinifera</i> , Robertson, South Africa
	LM23 = CBS 110835	AY179932	AY179898		<i>Vitis vinifera</i> , Robertson, South Africa
	LM34 = CBS 110708	AY179933	AY179899		<i>Vitis vinifera</i> , Paarl, South Africa
	LM74 = CBS 110709	AY179928	AY179894		<i>Vitis vinifera</i> , Paarl, South Africa
	LM119 = CBS 110834	AY179937	AY179903		<i>Vitis vinifera</i> , Paarl, South Africa
	LM466	AY179936	AY179902		<i>Vitis vinifera</i> , Paarl, South Africa
	LM440	AY179925	AY179891		<i>Vitis vinifera</i> , Wellington, South Africa
	CBS 246.91 ^d	AY179927	AY179893		<i>Vitis vinifera</i> , Wellington, South Africa
	CBS 100402	AY179934	AY179900		<i>Vitis vinifera</i> , Yugoslavia
	CBS 101006	AY179938	AY179904		<i>Vitis vinifera</i> , Italy
	CBS 101357	AY179939	AY179905		<i>Actinidia chinensis</i> , Italy
	CBS 100400	AY179942	AY179908		<i>Actinidia chinensis</i> , Italy
	CBS 100399	AY179941	AY179907		<i>Vitis vinifera</i> , Italy
	CBS 100398	AY179940	AY179906		<i>Vitis vinifera</i> , Italy
	LM24 = CBS 110833	AY179931	AY179897		<i>Vitis vinifera</i> , Italy
	LM44	AY179923	AY179889		<i>Vitis vinifera</i> , Paarl, South Africa
	LM52 = CBS 110832	AY179950	AY179916		<i>Vitis vinifera</i> , Paarl, South Africa
		AY179951	AY179917		<i>Vitis vinifera</i> , Paarl, South Africa

TABLE I. Continued

Species	Culture No.	GenBank No.				Host and location
		ITS	EF	SSU		
	LM75	AY179952	AY179919	AY179956	<i>Vitis vinifera</i> , Paarl, South Africa	
	LM83 = CBS 110829	AY179953	AY179921		<i>Vitis vinifera</i> , Paarl, South Africa	
	LM113	AY179954	AY179920		<i>Vitis vinifera</i> , Paarl, South Africa	
	LM115 = CBS 111016	AY179955	AY179918		<i>Vitis vinifera</i> , Paarl, South Africa	
	LM441	AY179924	AY179890		<i>Vitis vinifera</i> , Wellington, South Africa	
	LM443	AY179926	AY179892		<i>Vitis vinifera</i> , Wellington, South Africa	
	LM460	AY179922	AY179888		<i>Vitis vinifera</i> , Wellington, South Africa	
	CBS 213.31	AY179943	AY179909	AY179957	Unknown, Italy	
<i>Togninia novae-zealandiae</i>	UAMH 9589	AY179944	AY179910		<i>Cupressus macrocarpa</i> , New Zealand	
	UAMH 9590	AYA79445	AY179911	AY179958	<i>Pinus radiata</i> , New Zealand	
<i>Togninia fraxinopennsylvanica</i>	CBS 110212	AY179947	AY179913	AY179959	<i>Fraxinus pennsylvanica</i> , USA	
<i>Phaeacremonium</i> sp.	STE-U 3394	AY179946	AY179912		<i>Desmoschoenus spiralis</i> , New Zealand	

^a Designated mating type tester strains.

^b Mat1-1 mating type.

^c Mat1-2 mating type.

^d Ex-type culture of *Pm. aleophilum*.

DNA (SSU) gene for a subset of four isolates representing the different *Togninia* species and *Pm. aleophilum*. The cycling conditions consisted of an initial denaturation step of 94 C for 7 min, followed by 36 cycles of (1) denaturation (95 C for 45 s), (2) annealing (55 C for 60 s) and (3) elongation (72 C for 120 s), and finally a 2 min extension step at 72 C.

PCR products were analyzed by electrophoresis at 85 V for 30 min in a 0.8% (w/v) agarose gel in 0.5 × TAE buffer (0.4 M Tris, 0.05 M NaAc, and 0.01 M ethylene diamine tetraacetic acid [EDTA], pH 7.85) and visualized under UV light with a GeneGenius Gel Documentation and Analysis System (Syngene, Cambridge, United Kingdom) following ethidium bromide staining.

PCR products were purified according to the manufacturer's instructions using a commercial kit (Nucleospin Extract 2 in 1 Purification Kit, Machery-Nagel GmbH & Co., Germany). Sequencing reactions were carried out with ABI PRISM Big Dye Terminator version 3.0 Cycle Sequencing Ready Reaction Kit (PE Biosystems, Foster City, California), according to the manufacturer's recommendations, and were analyzed on an ABI Prism 3100 DNA Sequencer (Perkin-Elmer, Norwalk, Connecticut). Sequences were deposited at GenBank (TABLE I), and the alignment was deposited in TreeBase (ITS and EF-1α: SN1269–3614; SSU: SN1269–3617).

Phylogenetic analysis.—Raw sequence data were analyzed using EditView 1.0.1 (<http://www.appliedbiosystems.com>), and sequences were manually aligned by inserting gaps. Phylogenetic analyses were conducted using PAUP (Phylogenetic Analysis Using Parsimony) version 4.0b10 (Swofford 2000). Gaps were treated as a fifth character, and all characters were unordered and of equal weight. *Phialophora richardsiae* (Nannf.) Conant (CBS 270.33, GenBank ITS = AY179948, EF-1α = AY179914) and *Cercospora apii* Fresen. (CBS 119.25, GenBank ITS = AY179949, EF-1α = AY179915) were used as outgroups for both the EF-1α and ITS analyses. Six *Pm. aleophilum* isolates (LM 44, LM 52, LM 75, LM 83, LM 113, and LM 115) were excluded from the combined analyses. Their respective EF and ITS sequences were 100% similar to the sequences of LM 24, LM 441, LM 466, LM 443, LM 440, LM 463, LM 460, LM 34 and LM 5. Maximum-parsimony analysis was performed using the heuristic search option with a 1000 random-taxon additions and tree bisection and reconstruction (TBR) as the branch-swapping algorithm. Bootstrap support for the ITS and EF-1α analysis for internal branches was evaluated from 1000 heuristic search replicates and 1000 random taxon additions. Tree length, consistency index (CI), retention index (RI) and the rescaled consistency index (RC) values also were calculated. A partition homogeneity test in PAUP (Swofford 2000) was conducted to test the congruence between the ITS and EF-1α sequence datasets. Small subunit sequences were added to an alignment obtained from TreeBase (M911). Sequences representative of the different orders within the class Sordariomycetes, as well as the order Chaetothiales (Chaetothriomycetes), were retrieved from GenBank and added to the alignment. Neighbor-joining analyses of the SSU alignment (using uncorrected “p”,

Kimura-2-parameter and Jukes-Cantor substitution models) were done with PAUP version 4.0b10 (Swofford 2000). *Rhodosporidium toruloides* Banno and *Athelia bombacina* Pers. were used as outgroups for the neighbor-joining analysis.

RESULTS

Morphology.—In culture, *T. minima* strain CBS 213.31 produced a *Phaeoacremonium* anamorph similar to *Pm. aleophilum*, but distinct from the *Phaeoacremonium* anamorphs associated with *T. fraxinopennsylvanica* and *T. novae-zealandiae*. A detailed description of *T. minima*, based on the teleomorphs formed by isolates of *Pm. aleophilum* (TABLE I), as well as the material designated by Hausner et al (1992), is provided below:

- Togninia minima* (Tul. & C. Tul.) Berl., Icon. Fung. 3: 11. 1900. FIGS. 1–24
 ≡ *Calosphaeria minima* Tul. & C. Tul., Sel. Fung. Carpol. 2: 112, plate XIII:23–24. 1863.
 ≡ *Calosphaeria (Erostella) minima* (Tul. & C. Tul.) Sacc., Syll. Fung. 1: 101. 1882.
 ≡ *Erostella minima* (Tul. & C. Tul.) Traverso, Fl. Ital. Crypt. 1: 156. (1905) 1906.
 = *Calosphaeria alnicola* Ellis & Everh., Proc. Acad. Nat. Sci. Phila. 221. (1890) 1891.
 ≡ *Togninia alnicola* (Ellis & Everh.) Berl., Icon. Fung. 3: 10. 1900.
 = *Longoa paniculata* Curzi, Atti Ist. Bot. R. Univ. Pavia, Ser 3, 3: 204. 1927.

Anamorph. *Phaeoacremonium aleophilum* W. Gams, Crous, M.J. Wingf. & Mugnai, Mycologia 88: 791. 1996.

Mycelium consisting of branched, septate hyphae; hyphae occurring singly or in strands of up to 10, tuberculate (with warts to 1 μm) to verruculose, pale brown, becoming paler toward the conidiogenous region, 1.5–3 μm wide. *Chlamydoconidia* absent. *Perithecia* heterothallic, mostly aggregated, not valloid, sometimes solitary, mostly subepidermal also on the surface of the epidermis; perithecia subglobose, sometimes obpyriform, with a long cylindrical neck, (160–)250–285(–420) μm diam and basal part (200–)285–325(–400) μm tall. Wall consisting of two regions of *textura angularis*: outer region dark brown, cells smaller and more rounded than inner layer, approx. 8–10 cells thick (individual cells not visible further outward), 20–40 μm thick; inner region hyaline (centrum) to pale brown, 5–7 cells and 12–28 μm thick; surface covered with brown, septate hyphal appendages that become hyaline towards their tips (more abundant on older perithecia). *Perithecial necks* black, 1–3(–6) per perithecium, curved, verruculose, with apex often proliferating secondarily upon aging and then appearing nodulose; nodules


(–120 μm wide) also appearing lower down on the neck; necks 800–1800 (av. 1055) μm long, 35–130 (av. 69) μm wide at the base, and 20–60 (av. 41) μm wide at the apex, neck sometimes dividing into two near the apex. Multinecked perithecia often with a thin wall dividing the perithecial chamber. *Paraphyses* hyaline, septate, cylindrical, narrowing towards the tip, 45–125 (av. 83) μm long, 2–4 μm wide at the base and 1.5–2 μm at the apex, persistent. *Asci* arising in acropetal succession from sympodially proliferating ascogenous hyphae that appear spicate when mature, hyaline, clavate, with bluntly rounded apices and with sides straight or tapering towards the truncate or bluntly obtuse bases (17–)19–20(–27) \times 4–5 μm ; *apical complex* 0.5–1 μm thick, of indistinct structure, with a nonamyloid apical ring (negative in Melzer's reagent). *Ascogenous hyphae* hyaline, branched, smooth-walled, 2–3 μm wide (inflated bases –5 μm wide). *Ascospores* 1-celled, hyaline, oblong-ellipsoidal to allantoid with rounded ends, sometimes containing small guttules at the ends, (4–)4.5–5(–6.5) \times 1–2 μm (av. 5 \times 1.5 μm).

Conidiophores mostly micronematous, arising from aerial or submerged hyphae, erect, simple, frequently reduced to conidiogenous cells, rarely 1–2-septate, subcylindrical, pale brown, paler towards the tip, smooth to verruculose, straight to gently curved, 4–40 μm tall, 2–3 μm wide. *Conidiogenous cells* terminal or lateral, mostly monophialidic, subcylindrical to narrowly ellipsoidal, smooth to verruculose, subhyaline, 3–21 μm long, 1–2.5 μm wide at the base, 1–1.5 μm wide at the apex, with a terminal, inconspicuous, almost convergent, 0.5–1 μm long, 1 μm wide collarette. *Conidia* aggregating in slimy heads, hyaline, oblong-ellipsoidal to allantoid, when larger oblong to reniform, becoming 2-guttulate with age, (2.5–)3–4.5(–7) \times 1.5–2.5(–3) μm (av. 4 \times 2 μm).


Substrates and geographical distribution. In vascular bundles of vines and twigs of woody plants: *Vitis vinifera* (Argentina, Australia, Chile, France, Italy, New Zealand, South Africa, Spain, Turkey, U.S.A. and Yugoslavia), *Actinidia sinensis* (Italy), bark of a tropical rain forest tree (Papua New Guinea), *Olea europaea* (Italy) (Crous et al 1996, Larignon et al 1997, Ari 2000, Crous and Gams 2000, Armengol et al 2001, Groenewald et al 2001).

Cultural characteristics. See Crous et al (1996).


Type specimens. YUGOSLAVIA. On roots and stems of *Vitis vinifera*, 1990, *M. Muntañola-Cvetković* (CBS 246.91 dried specimen and ex-type culture of *Pm. aleophilum*). SOUTH AFRICA. WESTERN CAPE PROVINCE: Wellington and Paarl, respectively, stems of *Vitis vinifera*, 2001, *L. Mostert*, LM 463 (MAT1-1 = CBS 111015) \times LM 54 (MAT1-2 = CBS 110703),



FIGS. 1-7. *Togninia minima* and its anamorph *Phaeacremonium aleophilum*. 1. Perithecia. 2. Asci with spicate arrangement on ascogenous hyphae. 3. Paraphyses. 4. Ascospores. 5. Asci. 6. Conidiophores and conidiogenous cells. 7. Conidia. Bars = 10 μ m.



FIGS. 8–17. *Togninia minima*. 8. Perithecia on *Vitis* cane. 9. Perithecium with three necks. 10. Perithecium with globoid spore mass. 11. Vertical section through a perithecium. 12, 13. Asci with spicate arrangement on ascogenous hyphae. 14. Vertical section through a perithecium wall. 15. Asci and paraphyses. 16. Asci. 17. Ascospores. Bars = 40 μm in FIGS. 8–11, 10 μm in FIGS. 12–17.



FIGS. 18–24. *Phaeoacremonium aleophilum*, anamorph of *Togninia minima*. 18–21. Conidiophores and conidiogenous cells. 22. Conidia after 3 mo. 23. Conidia aggregated in globoid masses. 24. Conidia after 7 d. Bar = 10 μm .

[**epitype** of *T. minima*, designated here] dried specimen, herb. CBS 6580.

Notes. No information previously has been available regarding the anamorph of *T. minima*. The only culture currently available that previously has been identified as *T. minima* is CBS 213.31. It differs from freshly isolated strains of *Pm. aleophilum* in several morphological characters. *Phaeoacremonium aleophilum* isolates in general have buff (19"d) to honey (21"b) colonies, while colonies of CBS 213.31 are white to buff (19"d) with woolly mycelial tufts. The difference might be due to cultural degeneration. Conidiogenous cells (1.5 μm at the base) and conidia (1 μm wide) of CBS 213.31 were slightly narrower than those of *Pm. aleophilum* (2.5 and 2 μm , respectively).

Phaeoacremonium aleophilum differs from anamorphs of *T. fraxinopennsylvanica* and *T. novae-zealandiae*. Conidia of *T. fraxinopennsylvanica* and *T. novae-zealandiae* were up to 9 and 10.5 μm long, respectively, significantly longer than those of *Pm. aleophilum*, which did not exceed 7 μm . Also, the collarettes of *T. fraxinopennsylvanica* and *T. novae-*

zealandiae (1.0–1.8 μm) were longer than those of *Pm. aleophilum* (0.5–1 μm).

Matings.—Perithecia produced by crossing *Pm. aleophilum* isolates were contrasted with those described for *T. minima*, *T. fraxinopennsylvanica* and *T. novae-zealandiae* (TABLE II). No interspecies mating was observed, nor were any matings obtained with CBS 213.31. The teleomorph of *Pm. aleophilum* was identical, however, to that described by Hausner et al (1992) for *T. minima*. Furthermore, ascospores of *T. minima* are longer and perithecia are wider and have longer necks than those of *T. fraxinopennsylvanica* and *T. novae-zealandiae* (TABLE II).

Of the 21 isolates of *Pm. aleophilum* that were mated, 10 belonged to one mating type and 11 to the other (FIG. 25). The bi-allelic heterothallic mating system suggested by these results was verified with crosses among the F1 ascospore progeny. Of 20 single-spore isolates used from one perithecium, 10 grouped in one mating type and 10 in the other, thus agreeing with a 1:1 Mendelian segregation of mating type (FIG. 26).

TABLE II. Perithecial morphology of the various *Togninia* species studied

	Ascospore shape and dimensions (μm)	Ascus dimensions (μm)	Perithecial dimensions (μm)	Neck length (μm)	Reference
<i>T. minima</i>	Allantoid 5.0–6.5 \times 1–2	20–30 \times 4.5–6	225–490 (–525) diam		Tulasne and Tulasne (1863)
<i>T. minima</i>	3–5.5 \times 1–2	15–21 (–25) \times 3.5–5 (–6)	220–490 diam		Berlese (1900)
<i>T. minima</i>	Oblong-ellipsoidal to allantoid (4–)4.7–5 (–6.5) \times 1–2	(17–)19–20 (–27) \times 4–5	(160–)250–285 (–420) diam (200–)285–325 (–400) tall	800–1800	Present study
<i>T. fraxinopennsylvanica</i>	Cylindrical to slightly curved 4–5.2 \times 1.6–1.8 (–2)	16.5–21 \times 4–4.8	195–290 diam	400–900	Hausner et al (1992)
<i>T. novae-zealandiae</i>	Ellipsoidal to oblong-ellipsoidal, slightly rounded at the ends 3.8–5.6 \times (1.8–)2.2–2.6	15.5–24 \times 4.7–5.5 (–6.5)	175–265 diam	450–1300	Hausner et al (1992)

Phylogenetic analysis.—The combined alignment (ITS and EF-1 α) had a total length of 863 characters, of which 301 were constant, 329 were parsimony uninformative and 233 were parsimony informative. The result of the partition homogeneity test showed that the ITS and EF-1 α datasets were congruent ($P = 0.72$) and therefore could be combined. Parsimony analysis of the combined datasets, using a heuristic search with 1000 random taxa additions, resulted in 10 most-parsimonious trees (Tree length = 901 steps, CI = 0.937, RI = 0.903, RC = 0.846 and HI = 0.063). *Togninia minima* (CBS 213.31) grouped in the clade with all *Pm. aleophilum* sequences, as well as the ex-type strain of *Pm. aleophilum*. The analysis clearly delimits the three *Togninia* species with good bootstrap support (FIG. 27). Some variation was observed within the *Pm. aleophilum* clade, as two isolates (CBS 100400, 101357) formed a subclade with 100% bootstrap support within the *Pm. aleophilum* clade. These two isolates differed from the rest of the *Pm. aleophilum* isolates at 11 (for EF-1 α) and two (for ITS) nucleotide positions. Another group within the *Pm. aleophilum* clade also received significant support (87%). These sequences, however, differed from the other isolates in the *Pm. aleophilum* clade at only two nucleotide positions in the EF-1 α dataset. A total of 21 characters proved variable in the combined dataset (2.4%). The EF-1 α area was found to be more variable (15 nucleotides, 4.6%) than the ITS area (6 nucleotides, 1.1%).

In the phylogram of the SSU sequence data (FIG. 28), *Pm. aleophilum*, *Togninia minima* (CBS 213.31), *T. fraxinopennsylvanica* and *T. novae-zealandiae* grouped together, forming the Calosphaerales clade (Sordariomycetes) (100% bootstrap). *Magnaporthe grisea* (T.T. Hebert) M.E. Barr, as well as a teleomorph of *Harpophora* W. Gams (*Gaeumannomyces graminis* (Sacc.) Arx & D.L. Olivier, Magnaporthaceae, incertae sedis), formed a well-supported subclade (100% bootstrap) within the class Sordariomycetes. *Phialophora verrucosa* Medlar, anamorphic *Capronia semiimera* (Cand. & Sulmont) Unter. & F.A. Naveau clustered with *Capronia* spp. (Chaetothyriales) with 100% bootstrap support.

DISCUSSION

Togninia is accepted as being typified by *T. minima* (Clements and Shear 1931). This viewpoint has been followed by Hausner et al (1992), as explained in Holm (1992), and also is accepted by Barr (M. E. Barr pers comm, 20 Feb 2002), contrasting with her earlier view (Barr et al 1993) that *Togninia* was a synonym of *Pleurostoma* Tul. & C. Tul. A more complicated issue addressed by Hausner et al (1992) was

Isolate	LM44	LM45	LM46	LM47	LM48	LM49	LM50	LM51	LM53	LM54	LM56	LM58	LM61	LM65	LM76	LM77	LM78	LM458	LM463	LM467	LM468	
LM44	-																					
LM45	-	-																				
LM46	+	+	-																			
LM47	-	-	+	-																		
LM48	-	-	+	-	-																	
LM49	+	+	-	+	+	-																
LM50	-	-	+	-	-	+	-															
LM51	+	+	-	+	+	-	+	-														
LM53	-	-	+	-	-	+	-	+	-													
LM54	-	-	+	-	-	+	-	+	-	-												
LM56	+	+	-	+	+	-	+	-	-	+	-											
LM58	+	+	-	+	+	-	+	-	+	+	-	-										
LM61	-	-	+	-	-	+	-	+	-	-	+	+	-									
LM65	+	-	-	+	+	-	+	-	+	+	-	-	+	-								
LM76	+	+	-	+	+	-	+	-	+	+	-	-	+	-	-							
LM77	-	-	-	-	-	+	-	+	-	-	+	+	-	+	+	-						
LM78	+	-	-	+	+	-	+	-	+	+	-	-	+	-	-	+	-					
LM458	-	-	+	-	-	+	-	+	-	-	+	+	-	+	+	-	+	-				
LM463	+	+	-	+	+	-	-	-	-	+	-	-	+	-	-	+	-	-	-			
LM467	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	+	-	-	-		
LM468	-	+	-	+	+	-	+	-	+	+	-	-	+	-	-	+	-	+	-	-	-	-

FIG. 25. Schematic representation of a mating study with single conidial isolates of *Phaeacremonium aleophilum*. A (-) indicates that no perithecia formed, while (+) indicates formation of perithecia that exuded copious amounts of fertile ascospores.

whether the genus name *Togninia* had precedence over *Erostella*. *Erostella* first was described as subgenus of *Calosphaeria* Tul. & C. Tul., and subsequently was elevated to generic level as *Erostella* (Sacc.) Trav. (1905) and again as *Erostella* (Sacc.) Sacc. (1906). Both *Erostella* and *Togninia* have *Calosphaeria minima* Tul. & C. Tul. as lectotype. The key to resolving this mystery lies in the precise interpretation of the Latin description of *Togninia* provided by Berlese (1900), who placed 12 species and one variety in this genus. The decision of Clements and Shear (1931) to des-

ignate *T. minima* as lectotype of *Togninia* is not obviously supported by Berlese's text. Berlese (1900, p. 20), stated "*C.[alosphaeria] herbicola* E.[llis] et E.[verhart] id. *C. ambigua* Berl.[ese] est novi generis *Togninia* typus." Two interpretations of this text are possible. The first is that Berlese actually typified *Togninia* by *T. ambigua* Berl.; in this case the lectotypification of Clements and Shear (1931) should be rejected and *Erostella* can be resurrected for the taxa treated by Hausner et al (1992), with *Calosphaeria minima* as lectotype. The second interpretation,

Isolates	LM227	LM228	LM229	LM230	LM231	LM232	LM233	LM234	LM235	LM236	LM237	LM238	LM239	LM240	LM243	LM244	LM245	LM246	LM247	LM249	
LM227	-																				
LM228	+	-																			
LM229	-	+	-																		
LM230	-	+	-	-																	
LM231	+	-	+	+	-																
LM232	+	-	+	+	-	-															
LM233	+	-	+	-	-	-	-														
LM234	-	+	-	-	+	+	+	-													
LM235	+	-	+	-	-	-	-	-	-												
LM236	+	-	+	+	-	-	-	-	-	-											
LM237	-	+	-	-	+	+	+	-	-	+	-										
LM238	+	-	+	+	-	-	-	+	-	-	-	-									
LM239	-	-	-	+	-	-	-	-	-	-	+	-	-								
LM240	+	-	+	-	-	-	-	+	-	-	-	-	-	-							
LM243	+	-	+	+	-	-	-	-	-	-	+	-	-	-	-						
LM244	-	+	-	-	+	+	+	-	+	+	-	+	+	+	-						
LM245	-	+	-	-	+	+	+	-	+	+	-	+	+	+	+	-	-				
LM246	-	+	-	-	-	+	+	-	+	+	-	+	-	+	+	-	-	-			
LM247	-	+	-	-	+	+	+	-	+	+	-	+	+	+	+	-	-	-	-		
LM249	-	+	-	-	+	+	+	-	+	+	-	+	-	+	+	-	-	-	-	-	

FIG. 26. Schematic representation of crosses from single ascospores of *Togninia minima* obtained from a fertile mating (LM 54 × LM 463). A (-) indicates that no perithecia formed, while (+) indicates formation of perithecia that exuded copious amounts of fertile ascospores.




FIG. 27. One of 10 most-parsimonious trees obtained from heuristic searches of a combined alignment of the 5.8S rRNA gene and flanking ITS1 and ITS2 regions and the translation elongation factor 1-alpha gene (length = 901 steps, CI = 0.937, RI = 0.903, RC = 0.846 and HI = 0.063). Bootstrap support values (1000 repl.) above 65% are shown at the nodes. *Phialophora richardsiae* and *Cercospora apii* were used as outgroups.

which was explained by Holm (1992) and subsequently followed by Hausner et al (1992), is that Berlese did not consider *C. herbicola* to be synonymous with *C. ambigua*. Under the genus *Togninia*, Berlese (1900, p. 9) listed only *T. ambigua*, while he transferred *C. herbicola* to *Jattaea* on p 8 but *C. minima* to *Togninia* on p 11. The Latin “id.”, “idem”, means “the same as”, but it also has been translated as “ditto”, which is more appropriate here, as Berlese (1900) clearly did not treat *C. herbicola* as synonymous with *C. ambigua* but actually placed them in different genera. The species in this *Calosphaeria* complex are thus “novi generis *Togninia* typus”, meaning that they are of the *Togninia*-type (Holm

1992), and consequently the lectotypification of *Togninia* with *T. minima*, as proposed by Clements and Shear (1931), can be accepted.

Because no authentic material of *T. minima* exists, Hausner et al (1992) designated the original illustration (Tulasne and Tulasne 1863) as iconotype. *Togninia* thus is conceived as a genus that has solitary to clustered, globose perithecia with papillate to beak-like apices that can be smooth or ornamented. Asci are unitunicate with truncate bases and thickened apices, appearing in a spicate arrangement on ascogenous hyphae. Paraphyses are present, hyaline, septate, and ascospores are hyaline, aseptate, allantoid to ellipsoidal. Anamorphs are acremonium- to phi-




FIG. 28. Neighbor-joining phylogenetic tree obtained from partial 18S rDNA sequences. Bootstrap support values (1000 repl.) are shown above the nodes. *Taphrina deformans* and *Rhodosporidium toruloides* were used as outgroups. Sequences marked with an asterisk were taken from TreeBase (M911).

alophora-like (Hausner et al 1992). Hausner et al (1992) based their redescription of *T. minima* in part on two specimens, one collected from *Alnus* in the United States (N.A.F. 2514) and another from *Prunus pennsylvanica* collected in Canada (SSMF 725–7179). No cultures, however, were available for study. Clements and Shear (1931) treated *Longoa* Curzi as synonym of *Calosphaeria*, a genus that is heterogeneous (M. E. Barr pers comm). In her revision of the Calosphaerales, Barr (1985) did not treat *Longoa*, but based on its morphology Eriksson and Hawksworth (1986) considered the type species, *Longoa paniculata* (CBS 213.31, ex-type), to be a synonym of *T. minima*. This synonymy is supported by the origi-

nal description and molecular data obtained in this study.

A comparison of our fungus with the original descriptions of *Togninia* provided by Tulasne and Tulasne (1863) and Berlese (1900), as well as with the redescription of Hausner et al (1992), shows that our fungus clearly belongs in *Togninia*. Furthermore, its generic relationship with *T. fraxinopennsylvanica* and *T. novae-zealandiae* is confirmed via the tight cluster seen in 18S SSU sequence data (FIG. 28).

Before any teleomorph was known for *Phaeoacremonium*, the genus was considered to have affinities for the Magnaporthaceae (Dupont et al 1998). Members of this family generally are characterized by hav-

ing long, hairy necks and septate ascospores (Kirk et al 2001). The family is considered to be close to the Diaporthales (Winka and Eriksson 2000). The *Togninia* teleomorphs of *Phaeoacremonium* differ strongly from those of other Magnaporthaceous fungi. In addition, our 18S rDNA sequence analysis shows that the *Togninia* and *Phaeoacremonium* species investigated here form a distinct clade apart from the Diaporthales (FIG. 28), supporting their placement in the order Calosphaerales for reasons outlined by Barr (1985). Phylogenetic analyses of DNA sequence data have shown that *T. minima* forms a separate cluster with *T. fraxinopennsylvanica* and *T. novae-zealandiae*, as well as other species presently known in *Phaeoacremonium* (Groenewald et al 2001). *Togninia* teleomorphs also have been induced for several other *Phaeoacremonium* spp., which await further description once their mating strategies have been resolved.

Results from mating studies clearly have shown that *T. minima* has a biallelic heterothallic mating system. Such systems in fungi have been reviewed by Glass and Nelson (1994). Field isolates obtained from individual diseased vines could be induced to form the teleomorph in vitro on cane sections, indicating that compatible mating types co-occur in nature on the same vine. The conditions necessary for perithecial formation in the field remain unknown. The extent to which teleomorphs occur in the field also is unknown. The discovery of the teleomorph is important for the design and deployment of disease-control strategies, as well as for the overall understanding of the epidemiology of Petri disease.

ACKNOWLEDGMENTS

The authors acknowledge Winetech and the South African National Research Foundation (NRF) for financial support. Dr. Margaret E. Barr (B.C., Canada) is thanked for her comments on the status of the Calosphaerales, while Drs. James Reid and Georg Hausner (Univ. Manitoba, Canada) are thanked for their numerous letters, for providing copies of all their correspondence related to the *Togninia/Erostella* arguments and for taking the trouble to revive ex-type cultures of the species treated in their study.

LITERATURE CITED

- Ari ME. 2000. A general approach for esca disease in the vineyards of Turkey. *Phytopathol Mediterr* 39:35–37.
- Armengol J, Vicent A, Torné L, García-Figueres F, García-Jeménez J. 2001. Fungi associated with esca and grapevine declines in Spain: a three-year survey. *Phytopathol Mediterr* 40:S325–S329.
- Barr ME. 1985. Notes on the Calosphaerales. *Mycologia* 77: 549–565.
- , Rogers JD, Ju JM. 1993. Revisionary studies in the Calosphaerales. *Mycotaxon* 68:529–535.
- Berlese AN. 1900. *Icones fungorum omnium hucusque cognitorum*. Patavia 3:9–21.
- Carbone I, Kohn LM. 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91:553–556.
- Clements FE, Shear CL. 1931. *The genera of fungi*. New York: H.W. Wilson Co. 58 plates, p 496.
- Cottral E, Ridgeway H, Pascoe I, Edwards J, Taylor P. 2001. UP-PCR analysis of Australian isolates of *Phaeoconiella chlamydospora* and *Phaeoacremonium aleophilum*. *Phytopathol Mediterr* 40:S479–S486.
- Crous PW, Gams W. 2000. *Phaeoconiella chlamydospora* gen. et comb. nov., a causal organism of Petri grapevine decline and esca. *Phytopathol Mediterr* 39:112–118.
- , ——, Wingfield MJ, van Wyk PS. 1996. *Phaeoacremonium* gen. nov. associated with wilt and decline diseases of woody hosts and human infections. *Mycologia* 88:786–796.
- Dupont J, Laloui J, Roquebert MF. 1998. Partial ribosomal DNA sequences show an important divergence between *Phaeoacremonium* species isolated from *Vitis vinifera*. *Mycol Res* 102:631–637.
- Eriksson O, Hawksworth DL. 1986. Notes on Ascomycete Systematics. *Syst Ascom* 5:113–174.
- Gams W. 2000. *Phialophora* and some similar morphologically little-differentiated anamorphs of divergent ascomycetes. In: Seifert K, Gams W, Crous PW, Samuels GJ, eds. *Molecules, morphology and classification: towards monophyletic genera in the Ascomycetes*. *Stud Mycol* 45:187–199.
- Gatica M, Césari C, Magnin S, Dupont J. 2001. *Phaeoacremonium* species and *Phaeoconiella chlamydospora* in vines showing “hoja de malvón” and young vine decline symptoms in Argentina. *Phytopathol Mediterr* 40: S317–S324.
- Glass NL, Nelson MA. 1994. Mating-type genes in mycelial Ascomycetes. In: Wessels JGH, Meinhardt F, eds. *The mycota I: growth, differentiation and sexuality*. Berlin, Germany: Springer-Verlag. p 295–306.
- Groenewald M, Kang JC, Crous PW, Gams W. 2001. ITS and β -tubulin phylogeny of *Phaeoacremonium* and *Phaeoconiella* spp. *Mycol Res* 105:651–657.
- Hausner G, Eyjofsdottir GG, Reid J, Klassen GR. 1992. Two additional species of the genus *Togninia*. *Can J Bot* 70: 724–734.
- Holm L. 1992. On the typification of pyrenomycete generic names. *Syst Ascom* 2:29–30.
- Kirk PM, Cannon PF, David JC, Stalpers JA. 2001. *Ainsworth and Bisby's dictionary of the fungi*. 9th ed. Oxon, U.K.: CABI Publishing, CAB International. 655 p.
- Larignon P, Dubos B. 1997. Fungi associated with esca disease in grapevine. *Eur J Pl Pathol* 103:147–157.
- Lee SB, Taylor JW. 1990. Isolation of DNA from fungal mycelia and single spores. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. *PCR protocols: a guide to methods and applications*. New York: Academic Press. p 282–287.
- Mugnai L, Graniti A, Surico G. 1999. Esca (black measles)

- and brown wood-streaking: two old and elusive diseases of grapevines. *Pl Dis* 83:404–416.
- Péros JP, Jamaux-Despréaux I, Berger G. 2000. Population genetics of fungi associated with esca disease in French vineyards. *Phytopathol Mediterr* 39:150–155.
- Rayner RW. 1970. *A mycological colour chart*. Kew, Surrey: CMI and British Mycological Society. 17 sheets, 34 p.
- Rooney SN, Askalen A, Gubler WD. 2002. First report of the teleomorph of *Phaeoacremonium* spp., the cause of esca and decline of grapevines. *Phytopathol News* 36: 116.
- Scheck HJ, Vasquez SJ, Gubler WD. 1998. First report of three *Phaeoacremonium* spp. causing young grapevine decline in California. *Pl Dis* 82:590.
- Swofford DL. 2000. PAUP* 4.0: phylogenetic analysis using parsimony. Sunderland, Massachusetts: Sinauer Associates Inc.
- Tegli S. 2000. A hypothesis about the reproductive modes of *Phaeoacremonium aleophilum* and *Phaeomoniella chlamydospora*. *Phytopathol Mediterr* 39:289–298.
- , Bertelli E, Surico G. 2000a. Sequence analysis of ITS ribosomal DNA in five *Phaeoacremonium* species and development of a PCR-based assay for the detection of *P. chlamydosporum* and *P. aleophilum* in grapevine tissue. *Phytopathol Mediterr* 39:134–149.
- , Santilli E, Bertelli E, Surico G. 2000b. Genetic variation within *Phaeoacremonium aleophilum* and *P. chlamydosporum* in Italy. *Phytopathol Mediterr* 39:125–133.
- Tulasne LR, Tulasne C. 1863. *Selecta Fungorum Carpologia*. Paris. Vol. 2. Parisii: 34 plates, 319 p.
- White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. *PCR protocols: a guide to methods and applications*. New York: Academic Press. p 315–322.
- Winka K, Eriksson OE. 2000. *Papulosa amerospora* accommodated in a new family (Papulosaceae, Sordariomycetes, Ascomycota) inferred from morphological and molecular data. *Mycoscience* 41:97–103.