

Internet-Accessible DNA Sequence Database for Identifying *Fusaria* from Human and Animal Infections[∇]

Kerry O'Donnell,^{1*} Deanna A. Sutton,² Michael G. Rinaldi,² Brice A. J. Sarver,³
S. Arunmozhi Balajee,⁴ Hans-Josef Schroers,⁵ Richard C. Summerbell,⁶
Vincent A. R. G. Robert,⁷ Pedro W. Crous,⁷ Ning Zhang,⁸
Takayuki Aoki,⁹ Kyongyong Jung,¹⁰ Jongsun Park,¹⁰
Yong-Hwan Lee,¹⁰ Seogchan Kang,¹¹ Bongsoo Park,¹¹
and David M. Geiser¹¹

Bacterial Foodborne Pathogens and Mycology Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Peoria, Illinois¹; Department of Pathology, University of Texas Health Science Center, San Antonio, Texas²; Department of Biological Sciences, University of Idaho, Moscow, Idaho³; Centers for Disease Control and Prevention, Atlanta, Georgia⁴; Agricultural Institute of Slovenia, Ljubljana, Slovenia⁵; Sporometrics Inc., Toronto, Canada⁶; CBS-KNAW Fungal Biodiversity Center, Utrecht, Netherlands⁷; Department of Plant Biology and Pathology, Rutgers University, New Brunswick, New Jersey⁸; NIAS Genebank (MAFF), National Institute of Agrobiological Sciences, 2-1-2, Kannondai, Tsukuba, Ibaraki 305-8602, Japan⁹; Department of Agricultural Biotechnology, Seoul National University, Seoul 151-921, South Korea¹⁰; and Department of Plant Pathology, The Pennsylvania State University, University Park, Pennsylvania¹¹

Received 17 May 2010/Returned for modification 1 July 2010/Accepted 27 July 2010

Because less than one-third of clinically relevant *Fusaria* can be accurately identified to species level using phenotypic data (i.e., morphological species recognition), we constructed a three-locus DNA sequence database to facilitate molecular identification of the 69 *Fusarium* species associated with human or animal mycoses encountered in clinical microbiology laboratories. The database comprises partial sequences from three nuclear genes: translation elongation factor 1 α (*EF-1 α*), the largest subunit of RNA polymerase (*RPB1*), and the second largest subunit of RNA polymerase (*RPB2*). These three gene fragments can be amplified by PCR and sequenced using primers that are conserved across the phylogenetic breadth of *Fusarium*. Phylogenetic analyses of the combined data set reveal that, with the exception of two monotypic lineages, all clinically relevant *Fusaria* are nested in one of eight variously sized and strongly supported species complexes. The monophyletic lineages have been named informally to facilitate communication of an isolate's clade membership and genetic diversity. To identify isolates to the species included within the database, partial DNA sequence data from one or more of the three genes can be used as a BLAST query against the database which is Web accessible at FUSARIUM-ID (<http://isolate.fusariumdb.org>) and the Centraalbureau voor Schimmelfcultures (CBS-KNAW) Fungal Biodiversity Center (<http://www.cbs.knaw.nl/fusarium>). Alternatively, isolates can be identified via phylogenetic analysis by adding sequences of unknowns to the DNA sequence alignment, which can be downloaded from the two aforementioned websites. The utility of this database should increase significantly as members of the clinical microbiology community deposit in internationally accessible culture collections (e.g., CBS-KNAW or the *Fusarium* Research Center) cultures of novel mycosis-associated *Fusaria*, along with associated, corrected sequence chromatograms and data, so that the sequence results can be verified and isolates are made available for future study.

In addition to being the single most important genus of toxigenic phytopathogens (40), *Fusarium* (*Hypocreales*, *Ascomycota*) has emerged over the past 3 decades as one of the most important genera of filamentous fungi responsible for deeply invasive, opportunistic infections in humans (83). Clinically, fusarioses in immunocompetent patients typically present as superficial infections, such as onychomycosis and

trauma-associated keratitis, or locally invasive infections, such as sinusitis, catheter-associated peritonitis, pneumonia, or diabetic cellulitis (77). The 2005-2006 keratitis outbreaks within the United States and Asia, however, were unusual in that they were linked to the use of a novel soft contact lens cleaning solution, which was subsequently removed from the market (11). In contrast, immunocompromised or immunosuppressed patients who are persistently and profoundly neutropenic may acquire life-threatening angioinvasive, hematogenously disseminated fusarial infections associated with high morbidity and mortality rates (15). The high mortality of immunosuppressed patients is due in part to the broad resistance of most *Fusaria* to the spectrum of antifungals currently available (1, 4–7, 56); liposomal amphotericin

* Corresponding author. Mailing address: Bacterial Foodborne Pathogens and Mycology Research Unit, National Center for Agricultural Utilization Research, Agricultural Research Service, United States Department of Agriculture, 1815 North University Street, Peoria, IL 61604-3999. Phone: (309) 681-6383. Fax: (309) 681-6672. E-mail: kerry.odonnell@ars.usda.gov.

[∇] Published ahead of print on 4 August 2010.

B shows the greatest efficacy among the drugs currently in use (3, 17, 66).

A series of molecular phylogenetic studies has led to the important conceptual advance that morphological species recognition within *Fusarium* (22, 38, 47) greatly underestimates its species diversity (49, 50, 53, 54–57, 59, 70, 85). This finding is not too surprising, given that phenotypic methods for identifying fusaria rely on relatively few morphological and cultural characters (75). Based on an extensive literature review, Nucci and Anaissie (48) recently recorded 12 morphospecies associated with fusarial infections within the immunocompromised patient population. However, phylogenetic species recognition based on genealogical concordance of multilocus DNA sequence data (herein referred to as GCPSR) (79) has identified at least 69 clinically important *Fusarium* species (Table 1) (49, 54, 56, 57, 70, 85). Phylogenetic species in these studies were recognized if they received $\geq 70\%$ maximum parsimony (MP) bootstrap support (78) from the majority of the individual gene partitions and the combined data set and if their monophyly was not contradicted by analyses of any of the individual single-gene partitions.

Although GCPSR-based studies have revealed extensive cryptic speciation across the phylogenetic breadth of the genus and within other medically (9, 10, 33, 35, 65) and agriculturally important fungi (reviewed in references 21, 80, and 81), the level of cryptic speciation was especially pronounced within the *Fusarium solani* species complex (FSSC) (49, 56, 85) and *F. incarnatum*-*F. equiseti* species complexes (FIESC) (57). These two species complexes collectively harbor at least 75 species, including 41 associated with mycotic infection of humans and other animals. Multilocus DNA sequence data have proven to be essential for accurately circumscribing species boundaries within *Fusarium* and also have demonstrated utility in identifying epidemiologically important multilocus haplotypes, such as the widespread *F. oxysporum* clonal lineage (*F. oxysporum* species complex 3-a [FOSC 3-a], sequence types [ST] 33, 51, and 58) and FSSC 1-a and 2-d, which appear to be common in water systems (43, 54), including those of hospitals, where they pose a significant risk for nosocomial infections (2, 58).

Given the importance of fusaria to medicine, veterinary science, and agriculture, it is not surprising that diverse molecular methods for their identification have been published. The majority of these methods target the nuclear ribosomal internal transcribed spacer (ITS) region (30, 37, 67, 73) or domains D1 and D2 of the nuclear small-subunit ribosomal DNA (rDNA) (27, 28) as markers. Unfortunately, these methods were developed in reference to *Fusarium* morphospecies concepts, which greatly underestimate the species diversity reported herein based on GCPSR. Moreover, rDNA loci are too conserved to distinguish many closely related human pathogenic fusaria (8, 13, 54). Fortunately, recently published multilocus molecular phylogenetic studies of *Fusarium* have revealed that certain protein-encoding genes contain a wealth of phylogenetic signal (19, 53, 54, 56, 57, 70, 85). It is reasonable to assume that the genetic diversity of clinically and veterinarily relevant fusaria will continue to expand, whereas phenotypic methods will remain woefully inadequate for yielding accurate species-level identifications for over two-thirds of the fusaria encountered in the clinical laboratory. In response to this growing need for accurate species identification, the present study was initiated

with the aim of developing a comprehensive DNA sequence database that includes a representative of all presently known human/animal pathogenic *Fusarium* species identified previously using GCPSR.

Toward this end, a three-locus DNA sequence database for all known human opportunistic/pathogenic fusaria (i.e., 69 species) was developed to meet the following four objectives: (i) determine the utility of single- and multilocus DNA sequence data (*EF-1 α* , *RPB1*, and *RPB2*) for accurately identifying clinically important fusaria to species level, including partial sequence data from the DNA-directed RNA polymerase largest subunit (*RPB1*), which is used here for the first time for phylogenetic inference within *Fusarium*; (ii) investigate the phylogenetic diversity and evolutionary relationships of mycosis-associated fusaria; (iii) provide an Internet-accessible, three-locus database for accurately identifying and placing novel etiologic agents of fusarioses within a precise phylogenetic framework as they are encountered in the clinical microbiology laboratory; and (iv) archive a duplicate set of isolates at the CBS-KNAW in Europe and the ARS (NRRL) Culture Collection in the United States that is readily accessible to various research groups wanting to pursue further research on this topic. This *Fusarium* database, together with alignments and the corrected sequence chromatograms, will be incorporated into the FUSARIUM-ID database accessible via the Web at Pennsylvania State University (<http://isolate.fusariumdb.org>) and the Centraalbureau voor Schimmelcultures (CBS) Biodiversity Centre (<http://www.cbs.knaw.nl/fusarium>) to facilitate global identifications via the Internet and to promote cooperation and coordination in documenting and sharing the diversity and occurrence of clinically relevant fusaria.

MATERIALS AND METHODS

***Fusarium* isolates.** The 71 isolates included in this study comprise 69 phylogenetically distinct species (Table 1). Of these, 65 were cultured from human clinical or veterinary sources. Actual medical or veterinary case isolates were unavailable for three fusaria reported to cause infections in humans, so representative isolates of these three species (i.e., *F. napiforme*, *F. sporotrichioides*, and *F. lateritium*) from other sources were used as substitutes. With the exception of these three species, all of the other isolates have been characterized molecularly in published studies using partial DNA sequence data (see references in Table 1). The isolates included in this study are available upon request from the Agricultural Research Service (NRRL) Culture Collection (<http://nrri.ncaur.usda.gov/TheCollection/index.html>), National Center for Agricultural Utilization Research, Peoria, IL, and the CBS-KNAW, where they are stored cryogenically.

Molecular biology. Mycelia were grown in 300-ml Erlenmeyer flasks containing 100 ml of yeast-malt broth (20 g dextrose, 5 g peptone, 3 g yeast extract, and 3 g malt extract per liter; Difco, Detroit, MI) for 2 or 3 days on a rotary shaker at 100 rpm, harvested over a Büchner funnel, and then freeze-dried. Total genomic DNA was extracted from ~ 100 mg of freeze-dried mycelium using a cetyl trimethyl-ammonium bromide (CTAB; Sigma-Aldrich, St. Louis, MO) protocol as previously described (50). Portions of the translation elongation factor (*EF-1 α*) and DNA-directed RNA polymerase second largest subunit (*RPB2*) were selected based on their demonstrated utility in previous studies (54, 56, 57, 85). DNA-directed RNA polymerase subunit 1 (*RPB1*) was chosen based on published species-level studies from the Assembling the Fungal Tree of Life (AFTOL) project (18, 42). PCR and sequencing primers used for these three loci are provided in Table 2. Platinum *Taq* DNA polymerase (Invitrogen, Carlsbad, CA) was used in all PCRs which were conducted in an Applied Biosystems (ABI) 9700 thermocycler (Emeryville, CA) using published cycling parameters (50). Amplicons were size fractionated via gel electrophoresis in 1.5% agarose gels (Invitrogen) run in $1\times$ TAE buffer (69), stained with ethidium bromide and then photographed over a UV transilluminator. Prior to cycle sequencing, amplicons

TABLE 1. *Fusaria* subjected to DNA MLST

NRRL no.	Complex ^a	Species ^b	Equivalent no. ^c	Isolate source ^e	Geographic origin	Reference(s) or source
13604	GFSC	<i>F. napiforme</i>	CBS 748.97	Millet	Namibia	44
20423	FIESC	FIESC 4-a (<i>F. lacertarum</i>)	IMI 300797	Lizard skin	India	57, 74
20711	FDSC	<i>F. penzigii</i>	CBS 116508	Human eye	Sri Lanka	70
22608	FSSC	FSSC 20-a	UTHSC 93-1547	Human	Massachusetts	56
22611	FSSC	FSSC 14-a	UTHSC 93-2524	Human eye	Michigan	56
25197	FLSC	<i>F. cf. lateritium</i>	BBA 65687	<i>Bambusa vulgaris</i>	Venezuela	46
25229	GFSC	<i>F. thapsinum</i>	IMI 240460	Human mycetoma	Italy	6, 50
25378	FOSC	FOSC clade 3 ^d	IMI 214661	Human	Oklahoma	4, 51, 58
25387	FOSC	FOSC clade 2	ATCC 26225	Human toenail	New Zealand	4, 51, 58
25479	FSASC	<i>F. sporotrichioides</i>	CBS 447.67	<i>Pinus nigra</i> seed	Germany	60
25728	FCOSC	<i>F. concolor</i> ^e	CBS 463.91	Human	Germany	25
26360	FOSC	FOSC clade 1	FRC O-0755	Human eye	Tennessee	4, 51, 58
26421	GFSC	<i>F. nygamai</i>	CBS 140.95	Human	Egypt	36
28008	FSSC	FSSC 29-a	CDC B-4701	Human eye	Alabama	56
28009	FSSC	FSSC 15-a	CDC B-5543	Human eye	Texas	56
28029	FIESC	FIESC 3-b	CDC B-3335	Human eye	California	57
28541	FSSC	FSSC 26-a	UTHSC 98-1305	Human	Connecticut	56
28546	FSSC	FSSC 1-a	UTHSC 98-853	Human eye	Massachusetts	56
31158	FSSC	FSSC 18-a	MDA 1	Human	Texas	56
31169	FSSC	FSSC 25-a	MDA 12	Human	Texas	56
32309	FSSC	FSSC 12-d	UTHSC 00-1608	Human	Massachusetts	56
32434	FSSC	FSSC 16-b (<i>F. lichenicola</i>)	CBS 623.92	Human	Germany	56, 76
32437	FSSC	FSSC 28-a	CBS 109028	Human	Switzerland	56
32522	FIESC	FIESC 18-b	Loyola W-14182	Human diabetic cellulitis	Illinois	57
32755	FSSC	FSSC 9-a	FRC S-0452	Turtle	Florida	56
32864	FIESC	FIESC 17-a	FRC R-7245	Human	Texas	57
32865	FIESC	FIESC 21-b	FRC R-8480	Human endocarditis	Brazil	57
32866	FIESC	FIESC 23-a	FRC R-8822	Human cancer patient	Texas	57
32868	FIESC	FIESC 25-c	FRC R-8880	Human blood	Texas	57
32997	FIESC	FIESC 7-a	UTHSC 99-423	Human toenail	Colorado	57
34002	FIESC	FIESC 22-a	UTHSC 95-1545	Human ethmoid sinus	Texas	57
34003	FIESC	FIESC 20-a	UTHSC 95-28	Human sputum	Texas	57
34004	FIESC	FIESC 16-a	UTHSC 94-2581	Human BAL fluid	Texas	57
34005	FIESC	FIESC 24-a	UTHSC 94-2471	Human intravitreal fluid	Minnesota	57
34006	FIESC	FIESC 15-a	UTHSC 93-2692	Human eye	Texas	57
34016	FCSC	FCSC 2-a	UTHSC 98-2537	Human leg	Texas	57
34032	FIESC	FIESC 5-a	UTHSC 98-2172	Human abscess	Texas	57
34033	FSASC	<i>F. brachygibbosum</i>	UTHSC 97-99	Human foot cellulitis	Texas	57
34036	FTSC	FTSC <i>Fusarium</i> sp. 1	UTHSC 01-1965	Human ethmoid sinus	Colorado	57
36140	FDSC	<i>F. dimerum</i>	CBS 108944	Human blood	Netherlands	70
36147	FTSC	<i>F. acuminatum</i>	CBS 109232	Human bronchial secretion	Unknown	57
36160	FDSC	<i>F. delphinoides</i>	CBS 110140	Human eye	Florida	70
36185	FDSC	FDSC <i>Fusarium</i> sp. 5	CBS 110312	Human sinus	Washington	70
37393	FDSC	FDSC <i>Fusarium</i> sp. 2	FRC E-0105	Human eye	Sri Lanka	70
37625	FSSC	FSSC 27-a	CBS 518.82	Human	Netherlands	56
43433	FSSC	FSSC 2-a	CDC 2006011214	Human eye	Ohio	56
43441	FSSC	FSSC 3 + 4-a (<i>F. falciforme</i>)	CDC 2006743414	Human eye	Pennsylvania	56, 76
43467	FSSC	FSSC 8-a (<i>Fusarium</i> sp.) ^f	CDC 2006743430	Human eye	Louisiana	56
43468	FSSC	FSSC 5-a	CDC 2006743431	Human eye	Iowa	56
43489	FSSC	FSSC 6-a	CDC 2006743456	Human eye	Maryland	56
43498	FIESC	FIESC 8-b	CDC 2006743466	Human eye	Pennsylvania	57
43502	FSSC	FSSC 7-a	CDC 2006743470	Human eye	Tennessee	56
43608	GFSC	<i>F. verticillioides</i>	UTHSC 03-2552	Human peritoneal fluid	Minnesota	7, 75
43610	GFSC	<i>F. fujikuroi</i>	UTHSC 06-836	Human skin	Iowa	11
43617	GFSC	<i>F. proliferatum</i>	UTHSC 03-60	Human blood	Colorado	6, 50, 61, 75
43629	FCSC	FCSC 1-b	UTHSC 05-3200	Human blood	Utah	57
43631	FCSC	FCSC 3-a	UTHSC 05-2441	Human leg	Texas	57
43635	FIESC	FIESC 13-a	UTHSC 06-638	Horse	Nebraska	57
43636	FIESC	FIESC 14-c (<i>F. equiseti</i>)	UTHSC 06-170	Dog	Texas	57
43639	FIESC	FIESC 19-a	UTHSC 04-135	Manatee	Florida	57
43640	FIESC	FIESC 1-a	UTHSC 04-123	Dog nose	Texas	57
43641	FSASC	<i>F. armeniacum</i>	UTHSC 06-1377	Horse eye	Missouri	57
43694	FIESC	FIESC 6-a	CDC 2006743607	Human eye	Texas	57
44901	GFSC	<i>F. sacchari</i>	SSGH NC1	Human finger	Italy	6, 24
45999	FTSC	<i>F. flocciferum</i>	UTHSC 06-3449	Human scalp	California	57
46703	FSSC	FSSC 34-a	FMR 8281	Nematode	Spain	56
46707	FSSC	FSSC 35-a	FMR 8030	Human eye	Brazil	56

Continued on following page

TABLE 1—Continued

NRRL no.	Complex ^a	Species ^b	Equivalent no. ^c	Isolate source ^e	Geographic origin	Reference(s) or source
53131	GFSC	<i>F. ananatum</i>	SSGH VN	Human finger	Italy	31, 45
54126	GFSC	<i>F. acutatum</i>	FMR 8379	Human foot	Qatar	79
54147	FTSC	<i>FTSC Fusarium</i> sp. 2	CM 3913	Human pericardial fluid	Spain	1; this study
54158	GFSC	<i>F. subglutinans</i>	IUM 96-4102	Human blood	Italy	82

^a FCSC, *Fusarium chlamydosporum* species complex; FCOSC, *F. concolor* species complex; FDSC, *F. dimerum* species complex; FIESC, *Fusarium incarnatum-F. equiseti* species complex; FLSC, *F. lateritium* species complex; FOOSC, *F. oxysporum* species complex; FSASC, *F. sambucinum* species complex; FSSC, *F. solani* species complex; FTSC, *F. tricinctum* species complex; GFSC, *Gibberella (Fusarium) fujikuroi* species complex.

^b Arabic numerals identify species within species complexes; lowercase roman letters identify a unique haplotype within species (55, 56).

^c ATCC, American Type Culture Collection, Manassas, VA; BBA, Biologische Bundesanstalt für Land-und Forstwirtschaft, Institute für Mikrobiologie, Berlin, Germany; CBS-KNAW, Centraalbureau voor Schimmelcultures—Fungal Biodiversity Center, Utrecht, Netherlands; CDC, Centers for Disease Control and Prevention, Atlanta, GA; CM, Centro Nacional de Microbiología, Instituto de Salud Carlos III, Madrid, Spain; FMR, Facultat de Medicina i Ciències de la Salut, Reus, Spain; FRC, Fusarium Research Center, The Pennsylvania State University, State College, PA; IMI, CABI Biosciences, Egham, Surrey, England; IUM, Università degli Studi di Milano, Milan, Italy; Loyola, Loyola University, Maywood, IL; MDA, M. D. Anderson Cancer Center, Houston, TX; SSGH, Sesto San Giovanni Hospital, Milan, Italy; UTHSC, University of Texas Health Sciences Center, San Antonio, TX.

^d FOSC clades as reported by O'Donnell et al. (52).

^e Reported as *Fusarium polyphialidicum*, a later synonym of *F. concolor* (25).

^f FSSC 8 represents the homothallic species *Neocosmospora vasinfecta*, which produces an undescribed *Fusarium anamorph* (49).

^g BAL, bronchoalveolar lavage.

were purified using Montage₉₆ filter plates (Millipore Corp., Billerica, MA). Sequencing reactions were conducted in a 10-μl volume containing 2 μl of ABI BigDye Terminator, version 3.1, reaction mixture, 2 to 4 pmol of a sequencing primer, and approximately 50 ng of amplicon as previously described (50). After cycle sequencing, all reaction mixtures were cleaned up using an XTerminator purification kit and then run on an ABI 3730 48-capillary automated sequencer.

Phylogenetic analysis. Sequencher, version 4.9 (Gene Codes, Ann Arbor, MI), was used to edit and align raw ABI chromatograms, after which the *RPB1* and *RPB2* alignments were manually edited using TextPad, version 5.1.0 for Windows (Helios Software Solutions; Longridge, United Kingdom). Due to the presence of a number of length-variable indels within the three introns, sequences from the *EF-1α* partition were aligned automatically using MAFFT, version 6.0 (<http://align.bmr.kyushu-u.ac.jp/mafft/software/>), after which 92 ambiguously aligned intron nucleotide positions were excluded from the subsequent phylogenetic analyses. It is important to note that the entire region of *EF-1α* sequenced is

archived at the Fusarium-ID and CBS-KNAW databases. A conditional combination approach, which employed maximum parsimony bootstrap values of ≥70% as the threshold for topological discordance, indicated that the three individual partitions could be analyzed as a combined data set (Table 3). Phylogenetic relationships among the clinically relevant fusaria were inferred from the combined three-locus data set using unweighted MP implemented in PAUP, version 4.0b10 (78), and maximum likelihood (ML) employing GARLI, version 0.951 (86), as previously described (56). MrModeltest, version 3.8 (64), using the ModelTest server 1.0, identified the general-time-reversible model with a proportion of invariant sites and gamma-distributed rate heterogeneity (GTR+I+Γ) as the best-fit model of nucleotide substitution for the combined data set for the ML analyses. Searches for the shortest MP trees employed tree bisection and reconnection (TBR) branch swapping and 1,000 random sequence addition replicates. MP clade support was assessed by nonparametric bootstrapping, employing 1,000 pseudoreplicates of the data, 10 random addition sequences per rep-

TABLE 2. Primers used for PCR and DNA sequencing

Locus	Gene product	Length of sequence obtained (bp)	Primer		Use ^b		Reference(s) or source
			Designation	Sequence (5'–3') ^a	PCR	Sequencing	
<i>EF-1α</i>	Translation elongation factor 1α	639–683 ^c	EF1	ATGGGTAAGGARGACAAGAC	●		52
			EF2	GGARGTACCAGTSATCATG	●		52
			EF3	GTAAGGAGGASAAGACTCACC		●	56
			EF22T	AGGAACCCTTACCAGCTC		●	52
<i>RPB1</i>	RNA polymerase largest subunit	1,607	Fa	CAYAARGARTCYATGATGGGWC ^d	●		29
			G2R	GTCATYTGDDGTDGCDGGYTCDC	●		Benjamin Hall, unpublished data ^e
			R8	CAATGAGACCTTCTCGACCAGC	●		This study
			F5	ATGGGTATYGTCCAGGAYTC		●	This study
			F6	CTGCTGGTGGTATCATTACG		●	This study
			F7	CRACACAGAAGAGTTTGAAGG		●	This study
			F8	TTCTTCCACGCCATGGCTGGTCG		●	This study
R9	TCARGCCCATCGAGAGTTGTC	●		This study			
<i>RPB2</i>	RNA polymerase second largest subunit	1,700–1,742 ^f	5f2	GGGGWGAAYCAGAAGAAGGC	●	●	68
			7cr	CCCATRGCTTGYTTTRCCCAT	●	●	39
			7cf	ATGGGYAARCAAGCYATGGG	●	●	39
			11ar	GCRTGGATCTTRTCRTCSACC	●	●	39

^a D = A, G or T; R = A or G; S = C or G; W = A or T; Y = C or T.

^b ●, primer was used for indicated purpose. *RPB1* PCR primer G2R was designed by Benjamin Hall (<http://faculty.washington.edu/benhall/>).

^c The partial *EF-1α* sequence size range.

^d Fa is identical to *RPB1-DF2asc* (29), except that the 3'-most nucleotide was deleted.

^e <http://faculty.washington.edu/benhall/people.html>.

^f The size difference is due to 3-bp and 39-bp inserts in 5 × 7 *RPB2* fragment in members of the *Fusarium dimerum* species complex and the *Gibberella* clade, respectively.

TABLE 3. Tree statistics for the individual and combined partitions

Locus	No. of bp	PIC ^a	PIC/no. of bp	AUT ^b	No. of MPTs ^c	MPT length	CI ^d	RI ^e
<i>EF-1α</i>	632	284	0.45	43	>10,000	1,183	0.47	0.869
<i>RPB1</i>	1,607	603	0.38	52	24	2,590	0.34	0.851
<i>RPB2</i> 5 \times 7 ^g	882	379	0.43	26	13	1,475	0.43	0.879
<i>RPB2</i> 7 \times 11 ^h	860	265	0.31	30	618	1,272	0.35	0.83
<i>RPB2</i> 5 \times 7 and 7 \times 11	1,742	644	0.37	56	1,000	2,789	0.39	0.855
<i>RPB1</i> + <i>RPB2</i>	3,349	1,247	0.37	108	1	5,447	0.38	0.85
<i>EF-1α</i> + <i>RPB1</i> + <i>RPB2</i>	3,981	1,531	0.38	151	4	6,683	0.4	0.852

^a PIC, parsimony-informative character (i.e., shared derived nucleotide position or synapomorphy).

^b AUT, autapomorphy or a derived character unique to a particular taxon (i.e., not parsimony informative).

^c MPT, most parsimonious tree (i.e., shortest tree inferred from the DNA sequence data).

^d CI, consistency index. This index provides a metric of how much noise is in a data set. The CI is calculated by dividing the minimum possible number of steps by the observed number of steps.

^e RI, retention index. Like the CI, this metric provides a measure of how much noise is in a data set. Unlike the CI, the RI measures the amount of parsimony-informative characters from a data set that are reflected in the phylogenetic tree.

^f Ninety-two ambiguously aligned positions were excluded from the *EF-1 α* partition.

^g Region PCR amplified by primers 5f2 and 7cr (Table 2).

^h Region PCR amplified by primers 7cf and 11ar (Table 2).

licate, and TBR branch swapping. Nonparametric ML bootstrapping was conducted with a 2.6-GHz MacBook Pro, using 5,000 generations without improving the topology parameter and 1,000 ML pseudoreplicates of the data.

Nucleotide sequence accession numbers. DNA sequences have been deposited in GenBank under accession numbers HM347114 to HM347221.

RESULTS AND DISCUSSION

The primary objective of this study was to develop a Web-accessible three-locus DNA sequence database to facilitate identification of fusaria associated with human and animal infections. Additionally, the utility of single- and multilocus DNA sequence data for accurately identifying clinically important fusaria and a duplicate set of isolates at the CBS-KNAW in Europe and the ARS (NRRL) Culture Collection in the United States is described.

Phylogenetic relationships and identification of human pathogenic fusaria. The database was populated with aligned partial sequences from the nuclear genes *RPB1* (1,607 sites), *RPB2* (1,742 sites), and *EF-1 α* (632 sites) from 71 isolates representing 69 fusariosis-associated species reported in the literature (Table 1). Sequences of *F. sporotrichioides* and *F. lateritium* were included in the database, with the caveat that the single reports of these species causing infections in humans need to be verified. Although *F. napiforme* has clearly been shown to cause a human mycotic infection (44), we consider

reports of *F. sporotrichioides* (60) and *F. lateritium* (46) as etiological agents of fusarioses to be tentative because these identifications were not supported by molecular data and because these isolates were unavailable for further study. The morphological concepts of both species are known to comprise multiple phylogenetic species (20; K. O'Donnell, unpublished data). If identification of haplotypes within a species is required, detailed information on additional loci to sequence has previously been published (34, 51, 56, 57, 71, 85). NEXUS files with a PAUP block used to develop the FSSC, FIESC, *F. chlamyosporum* species complex (FCSC), *F. dimerum* species complex (FDSC), and *Gibberella fujikuroi* species complex (GFSC) can be downloaded from the Internet-accessible *Fusarium* database sites cited above or accessed via the dedicated BLAST servers.

PCR primers Fa and G2R, which were designed for higher-level phylogenetics as part of the AFTOL project (Table 2), successfully amplified an 1,894-bp fragment from the *RPB1* D-to-G region in all of the fusaria included in the database except for *F. cf. lateritium* NRRL 25197. The Fa and G2R primer sites in this isolate, however, appear to be conserved, based on DNA sequence analysis of overlapping fragments obtained using Fa/R8 and F7/R9 as PCR primers (Fig. 1). Because DNA sequence data from the *RPB1* locus had not been used previously for *Fusarium* phylogenetics, the design of

RPB1 primer map




FIG. 1. Map of the RNA polymerase largest-subunit (*RPB1*) locus. Location and orientation of PCR and sequencing primers are indicated by half-arrows. PCR primers Fa and G2R (Table 2) successfully amplified an 1,894-bp fragment in all of the isolates except for NRRL 25197 *Fusarium cf. lateritium*, even though the PCR primer sites appear to be conserved in this isolate. Therefore, the Fa \times G2R region was amplified in this isolate as two overlapping fragments using the PCR primer pairs Fa/R8 (1,127 bp) and F7/R9 (2,217 bp). Primers F5 and G2R flank the 1,607-bp *RPB1* D-to-G region that was sequenced and analyzed. Sequences for 65 of the 71 isolates were generated using the F5, F7, and F8 sequencing primers. See Fig. 1 in Matheny et al. (41) for a detailed map of the entire *RPB1* locus showing the position of the D-to-G region.

internal sequencing primers was accomplished by downloading sequences of this gene from the three sequenced fusarial genomes (*F. graminearum*, *F. oxysporum*, and *F. verticillioides*) at the Broad Institute of MIT and Harvard University (http://www.broadinstitute.org/annotation/genome/fusarium_group/MultiHome.html) and the genome of *F. solani* f. sp. *pisi* [*Nectria haematococca* from the Department of Energy Joint Genome Institute (JGI) (<http://genome.jgi-psf.org/Necha2/Necha2.home.html>)]. As reported for other filamentous ascomycetes (26), the *RPB1* D-to-G region in *Fusarium* was entirely exonic and free of indels. Alignment of the four *RPB1* sequences facilitated the design of five conserved internal sequencing primers (Fig. 1). However, only three were needed (i.e., F5, F7, and F8) to obtain reliable sequence coverage of 66 of the 71 fusaria included in this study. Two additional primers (i.e., F6 and Fa) were required to completely sequence the D-to-G region in members of the *F. dimerum* species complex (FDSC) and *F. fujikuroi* NRRL 43610. Alignment of the *RPB2* region between primers F5 and F7 required the insertion of two indels 3 and 39 bps in length to accommodate 1 and 13 additional codons, respectively, within members of the FDSC and the *Gibberella* clade. By way of contrast, due to the presence of three length-variable introns, we employed the software program MAFFT (<http://align.bmr.kyushu-u.ac.jp/mafft/software/>) to align the *EF-1 α* region. To obtain an alignment of the *EF-1 α* region that reflected positional homology, 92 intron nucleotide positions were coded as ambiguously aligned and excluded from all subsequent phylogenetic analyses.

The three-locus data set totaled 3,981 bp of aligned DNA sequence data, including 1,531 parsimony-informative nucleotide positions. Summary sequence and tree statistics for the individual and combined data sets indicated that the three loci contained very similar levels of parsimony-informative characters (PIC) per bp of sequence (Table 3). The best ML tree, based on 10 independent analyses of the concatenated data set, yielded a log likelihood of $-38,129.37$ (Fig. 2). The four most-parsimonious trees were 6,683 steps in length and differed only in minor rearrangements of four closely related phylogenetic species within the FIESC (19, 22–24) and members of the FOOSC (Table 3). Because the root position of the tree is unknown (54), the trees were midpoint rooted. Irrespective of whether trees were midpoint rooted or rooted using sequences of the FDSC or FSSC as a sister to the ingroup, the root always joined the tree with *F. cf. lateritium* NRRL 25197 forming the most basal divergence within a strongly supported (100% bootstrap support [BS]) *Gibberella* clade (Fig. 2). ML and MP phylogenetic analyses of the concatenated data set recovered trees that were highly concordant topologically (Fig. 2; only the best ML tree is shown). Evolutionary relationships among the six informally named species complexes within the *Gibberella* clade were fully resolved by ML bootstrapping. ML and MP bootstrapping recovered similar levels of clade support, with two exceptions. One of these involved the monophyly of the GFSC and its three biogeographically structured subclades (50). Although the GFSC and its three subclades were strongly supported by ML bootstrapping (Fig. 2), as were the previously inferred (American (Asian, African)) evolutionary relationships of the subclades (50), only the American clade received strong MP bootstrap support. In the second exception, the *F. tricinctum* species complex (FTSC) received moderate support

(79% BS) as a sister to the ((FCSC, *F. sambucinum* species complex [FSAMSC]) FIESC) clade in the ML analysis, but this relationship was not supported by MP bootstrapping (51% BS). Close to three-quarters of the medically relevant *Fusarium* species were nested within the following three species complexes: FSSC ($n = 21$), FIESC ($n = 20$), and GFSC ($n = 10$). Of these, members of the FSSC are by far the most important, accounting for approximately 50 to 60% of all fusarioses worldwide (5, 56, 85).

Web-based identification of human pathogenic fusaria. The database described in this paper is accessible via the Web in two forms with different features, either of which can be used for routine identification, and housed and maintained at Pennsylvania State University (<http://isolate.fusariumdb.org>) and at the CBS Fungal Biodiversity Center (<http://www.cbs.knaw.nl/fusarium>). FUSARIUM-ID was originally set up in 2004 (19) as the first dedicated website for the molecular identification of fusaria using a partial *EF-1 α* gene sequence of an unknown as the query to BLAST the database. Construction of the FUSARIUM-ID database was motivated in part to ensure that researchers could use sequence data to make connections between their isolates of interest and sequence-characterized isolates available in public culture collections. This is essential because queries of GenBank (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) consistently recovered *Fusarium* sequences that were incorrectly identified (discussed in reference 63). For the latter reason, researchers who choose to access the data generated in the present study via GenBank are advised to look at the top sequences sorted by maximum identity (“Max ident”) producing significant alignments, to make sure that the organism name is used consistently. FUSARIUM-ID has been updated with new data search and visualization tools (S. Kang, unpublished data) and currently archives multilocus DNA sequence data from selected species, including the broad spectrum of medically important fusaria, and sequence data from most previously published studies will be uploaded into FUSARIUM-ID. Expansion of FUSARIUM-ID should greatly facilitate accurate species identifications, especially in light of the fact that at least two-thirds (46/69) of the human pathogenic species cannot be identified currently using morphological data.

Using the FUSARIUM-ID database. A Web-accessible user’s guide to FUSARIUM-ID can be found at the following link: <http://isolate.fusariumdb.org/guide.php>. The updated FUSARIUM-ID database discussed here can be queried via the BLAST feature using sequence data or by other information associated with isolates, including host or substrate of origin, geographic origin, and accession numbers from other culture collections. We recommend using one of the three loci highlighted in this paper (*EF-1 α* , *RPB1*, or *RPB2*) first because of their relatively complete coverage of human pathogenic fusaria.

Also available are the DNA sequence alignments used in the various published multilocus phylogenetic and identification studies that are the basis for this database (49–59, 70, 85). Additional tools for manipulating DNA sequence data and for visualization of geographic distribution of isolates are available, as they are for the *Phytophthora* plant pathogen database (62). The Biolomics software package, utilized in the CBS strain database (<http://www.cbs.knaw.nl/fungi/BioLOMICS.aspx?searchopt=4>), provides a wide array of additional tools,




FIG. 2. Best maximum likelihood tree inferred from the combined three-locus data set for 71 isolates representing 69 medically and veterinarily important *Fusarium* species. Because the branching order of the two most basal lineages, the *F. solani* and *F. dimerum* species complexes (FSSC and FDSC), was unresolved in more inclusive analyses (54), the phylogram was midpoint rooted. The *Gibberella* clade contains the six most derived, clinically relevant species complexes. Species and their multilocus haplotypes are identified by Arabic numbers and lowercase Roman letters, respectively, for members of the *Fusarium incarnatum-F. equiseti* species complex (FIESC), the *F. chlamydosporum* species complex (FCSC), and the FSSC as previously reported (56, 57). Numbers in parentheses by the three *F. oxysporum* species complex (FOSC) isolates refer to clades as reported by O'Donnell et al. (50). Note that Latin binomials can be applied with confidence to only 23 of the 69 species. *Fusarium sporotrichioides* and *F. cf. lateritium* are highlighted in gray to indicate that reports of these species causing human infections need to be confirmed. Numbers above internodes represent ML bootstrap values based on 1,000 pseudoreplicates of the data. MP bootstrap values are indicated below internodes only when they differed by $\geq 5\%$ of the MP value. Af, African subclade; Am, American subclade; As, Asian subclade; FSAMSC, *F. sambucinum* species complex; FTSC, *F. tricinctum* species complex; and GFSC, *Gibberella fujikuroi* species complex.

including simultaneous searches with multiple loci utilizing the “multiple sequences” tool and searches against all of the accession numbers within the entire CBS fungal culture collection.

Interpreting the results. We anticipate that clinical microbiologists with access to DNA sequencing technology will utilize this database for identification of isolates to the species level, often using a single DNA marker (generally *EF-1 α* , *RPB1*, or *RPB2*) in doing so. In the context of other information, data from a single locus is often but not always sufficient to provide a reasonably high-confidence identification. The utility of a sequence match result depends on a variety of factors, including the quality of the sequence being used as a query, the degree to which the diversity of fusaria are represented in the database, and the various levels of known and actual DNA sequence diversity within species. All results must be interpreted with care, and precise conclusions may require additional data and analyses, including phylogenetic analysis.

Prior to conducting BLAST searches, it is essential that the sequences be edited using a software program such as DNA Chromatogram Explorer Lite (freeware available from HeracleSoftware, Lilienthal, Germany) or Sequencher (Gene Codes, Ann Arbor, MI) to trim low-quality ends and reconcile any differences between overlapping sequences. Whether using a partial *EF-1 α* , *RPB1*, or *RPB2* gene sequence as the query, an exact match to one of the human pathogenic species in the FUSARIUM-ID database generally can be interpreted as definite species-level identification. It is important for researchers to be aware, however, that several plant-pathogenic *Fusarium* species, including those that cause economically devastating diseases, such as fusarium head blight (FHB) of cereals (FSAMSC) (53, 59, 84) and soybean sudden death syndrome (SDS) (FSSC) (55) cannot be distinguished using DNA sequences of the three genes included in FUSARIUM-ID. For this reason, it is prudent to check the top sequences producing significant alignments to make sure that two or more species do not share the same allele. In addition to the FHB and SDS fusaria noted above, this problem may be encountered with a small number of clinically relevant species within the FSSC and FIESC. In anticipation of this problem, published sequence data comprising the three-locus typing schemes (*EF-1 α* , *RPB2*, and ITS+large subunit [LSU] rDNA) for these two complexes (56, 57) have been incorporated into the FUSARIUM-ID database. It should be noted that, compared to the partial *EF-1 α* , *RPB1*, and *RPB2* gene sequences, the ITS+LSU rDNA possesses relatively little phylogenetic signal within *Fusarium* (8, 54). Although the ITS rDNA region has been adopted widely by the fungal community as the universal locus for DNA barcoding of fungi (72), use of this marker within *Fusarium* for inferring evolutionary relationships is complicated by the presence of ITS2 paralogs (origin by gene duplication) or xenologs (origin by horizontal gene transfer) whose phylogenetic distribution does not track with the species phylogeny (50). Similarly, the discovery of highly divergent paralogs and low sequence divergence among orthologs of the mitochondrial cytochrome oxidase 1 gene in *Fusarium* (23), a locus widely promoted for barcoding diverse organisms (<http://www.barcoding.si.edu/DNABarCoding.htm>), indicates potential

problems in using this locus for phylogenetics and identification of fusaria.

A single-locus sequence query to the database may provide exact matches to isolates of one or more multilocus sequence types (MLSTs) defined in previous studies. Users are reminded that such an exact match reflects only the single locus used as a query; their isolate of interest may differ from the MLST matches at other loci and thus not fit into that MLST as heretofore defined. As discussed previously (19), when a query sequence does not perfectly match anything in the database, the unknown may represent a novel allele of a species in the database or a novel *Fusarium* species. For the majority of the queries, it is reasonable to assume that the unknown sequence represents a novel allele of a species represented in the database, given the database's dense taxon sampling. Assuming that this is the case, then the top match or matches should confirm that the variant allele is nested within a species previously characterized by GCPSR. However, when the unknown appears to be closely related to more than one species in the database, we recommend that additional sequence data be generated to take advantage of the wealth of multilocus DNA sequence data generated in published GCPSR-based studies of the FDSC (70), FSSC (49, 56, 85), FOCS (51, 58), GFSC (50), FCSC, and FIESC (57). Though representatives of the FTSC and FSAMSC are included in the current database, they are very rarely encountered as etiologic agents of fusarioses (57). If the BLAST results indicate that the query sequence represents a novel species not currently represented in FUSARIUM-ID, then the test isolate's genealogical exclusivity should be evaluated via GCPSR analyses, using the appropriate multilocus typing scheme and including two or more isolates, if available, to assess their monophyly via bootstrapping. In practice, we recognize such isolates as phylogenetically distinct species only if they are resolved as reciprocally monophyletic in the majority of the bootstrapped individual gene partitions, as well as in the combined data set, and their monophyly is not contradicted by bootstrapping of any individual partition (14, 56, 57).

While some phylogenetic species are known to possess little or no allelic variation within the major diagnostic markers employed, others are far more variable. Isolates of *F. proliferatum*, for example, may differ by as much as 2.1% at the *EF-1 α* locus and 1.7% at the *RPB2* locus (D. M. Geiser and K. O'Donnell, unpublished data). In most cases, a moderately divergent, single-locus best match (e.g., 96 to 98% identity at the *EF-1 α* locus) would likely represent a species that is not represented in the database, but data from additional loci and phylogenetic analyses would be necessary to determine that.

It is worth mentioning that the current version of FUSARIUM-ID possesses notable similarities and differences with TrichoKEY (<http://www.isth.info/tools/molkey/index.php>), a Web-accessible site dedicated to identification of *Trichoderma* spp. (16). Like FUSARIUM-ID, TrichoKEY provides a BLAST function to identify unknowns using DNA sequence data. FUSARIUM-ID has been updated to incorporate two useful features of TrichoKEY: (i) BLAST queries using three-locus DNA sequence data and (ii) the ability to download sequence alignments for subsequent phylogenetic analyses. These two databases differ primarily in that ITS rDNA has been reported to be more useful than partial *EF-1 α* and *RPB2* sequences for identifications within *Trichoderma* (12, 32),

whereas *EF-1 α* , *RPB1*, and *RPB2* appear to contain roughly equal levels of phylogenetic signal useful for species-level identifications within *Fusarium*. We also point out that the 5-to-7 region of *RPB2* often can be used alone, without including sequence data from the 7-to-11 region, to identify an unknown to the species level.

The utility of the fusariosis-associated portion of the FUSARIUM-ID data set is expected to grow as new validated sequences/sequence chromatograms and cultures are accessioned, especially as researchers and journals recognize the necessity for molecularly based identifications of fusarial pathogens. The Web-accessible FUSARIUM-ID database and the CBS database will continue to be updated with phylogenetically diverse partial *EF-1 α* , *RPB1*, and *RPB2* sequences, thereby enabling researchers to accurately identify virtually all unknowns to the species level, as well as allowing them to precisely place novel pathogens within a robust phylogenetic framework. Molecular phylogenetic clarification of human opportunistic fusarial species limits represents a significant conceptual and technological advance, which should help facilitate the long-term goals of epidemiologic studies directed at identifying the spectrum of etiologic agents and their environmental reservoirs, especially among hospitalized patients at greatest risk for acquiring nosocomial infections. Through the elucidation of species boundaries in the human-pathogenic fusaria, it should be possible to develop DNA sequence-independent methods for their rapid identification, such as microsphere (54, 55, 84) and oligonucleotide arrays (30).

ACKNOWLEDGMENTS

Special thanks are due to Stacy Sink and Jean H. Juba for excellent technical assistance, Nathane Orwig for generating the DNA sequences in NCAUR's DNA core facility, and the culture collections and individuals who supplied isolates used in this study.

The mention of trade products or firm names does not imply that they are recommended by the U.S. Department of Agriculture over similar products or other firms not mentioned. In addition, the findings and conclusions in this article are those of the author(s) and do not necessarily represent the views of the CDC.

REFERENCES

- Alastruey-Izquierdo, A., M. Cuenca-Estrella, A. Monzón, E. Mellado, and J. L. Rodríguez-Tudela. 2008. Antifungal susceptibility profile of clinical *Fusarium* spp. isolates identified by molecular methods. *J. Antimicrob. Chemother.* **61**:805–809.
- Anaisie, E. J., R. T. Kuchar, J. H. Rex, A. Francesconi, M. Kasai, F.-M. C. Müller, M. Lozano-Chiu, R. C. Summerbell, M. C. Dignani, S. J. Chanock, and T. J. Walsh. 2001. Fusariosis associated with pathogenic *Fusarium* species colonization of a hospital water system: a new paradigm for the epidemiology of opportunistic mold infections. *Clin. Infect. Dis.* **33**:1871–1878.
- Arikan, S., M. Lozano-Chiu, V. Paetznick, S. Nangia, and J. H. Rex. 1999. Microdilution susceptibility testing of amphotericin B, itraconazole, and voriconazole against clinical isolates of *Aspergillus* and *Fusarium* species. *J. Clin. Microbiol.* **37**:3946–3951.
- Azor, M., J. Cano, J. Gené, and J. Guarro. 2009. High genetic diversity and poor *in vitro* response to antifungals of clinical strains of *Fusarium oxysporum*. *J. Antimicrob. Chemother.* **63**:1152–1155.
- Azor, M., J. Gené, J. Cano, and J. Guarro. 2007. Universal *in vitro* antifungal resistance of genetic clades of the *Fusarium solani* species complex. *Antimicrob. Agents Chemother.* **51**:1500–1503.
- Azor, M., J. Gené, J. Cano, P. Manikandan, N. Venkatapathy, and J. Guarro. 2009. Less-frequent *Fusarium* species of clinical interest: correlation between morphological and molecular identification and antifungal susceptibility. *J. Clin. Microbiol.* **47**:1463–1468.
- Azor, M., J. Gené, J. Cano, D. A. Sutton, A. W. Fothergill, M. G. Rinaldi, and J. Guarro. 2008. *In vitro* antifungal susceptibility and molecular characterization of clinical isolates of *Fusarium verticillioides* (*Fusarium moniliforme*) and *Fusarium thapsinum*. *Antimicrob. Agents Chemother.* **52**:2228–2231.
- Balajee, S. A., A. M. Borman, M. E. Brandt, J. Cano, M. Cuenca-Estrella, E. Dannaoui, J. Guarro, G. Haase, C. C. Kibbler, W. Meyer, K. O'Donnell, C. A. Petti, J. L. Rodríguez-Tudela, D. Sutton, A. Velegraki, and B. L. Wickes. 2009. Sequence-based identification of *Aspergillus*, *Fusarium*, and *Mucorales* species in the clinical mycology laboratory: where are we and where should we go from here? *J. Clin. Microbiol.* **47**:877–884.
- Balajee, S. A., J. Houbraken, P. E. Verweij, S.-B. Hong, T. Yaguchi, J. Varga, and R. A. Samson. 2007. *Aspergillus* species identification in the clinical setting. *Stud. Mycol.* **59**:39–46.
- Bovers, M., F. Hagen, E. E. Kuramae, and T. Boekhout. 2008. Six monophyletic lineages identified within *Cryptococcus neoformans* and *Cryptococcus gattii* by multi-locus sequence typing. *Fungal Genet. Biol.* **45**:400–421.
- Chang, D. C., G. B. Grant, K. O'Donnell, K. A. Wannemuehler, J. Noble-Wang, C. Y. Rao, L. M. Jacobson, C. S. Crowell, R. S. Sneed, F. M. T. Lewis, J. K. Schaffzin, M. A. Kainer, C. A. Genese, E. C. Alfonso, D. B. Jones, A. Srinivasan, S. K. Fridkin, and B. J. Park. 2006. A multistate outbreak of *Fusarium* keratitis associated with use of a contact lens solution. *JAMA* **296**:953–963.
- Chaverri, P., and G. J. Samuels. 2004. *Hypocrea/Trichoderma* (Ascomycota, Hypocreales, Hypocreaceae): species with green ascospores. *Stud. Mycol.* **48**:1–116.
- Clinical and Laboratory Standards Institute. 2008. Interpretive criteria for microorganism identification by DNA target sequencing; proposed guideline. CLSI document MM18-A. Clinical and Laboratory Standards Institute, Wayne, PA.
- Dettman, J. R., D. J. Jacobson, and J. W. Taylor. 2003. A multilocus genealogical approach to phylogenetic species recognition in the model eukaryote *Neurospora*. *Evolution* **57**:2703–2720.
- Dignani, M. C., and E. Anaissie. 2004. Human fusariosis. *Clin. Microbiol. Infect.* **10**(Suppl. 1):67–75.
- Druzhinina, I. S., A. G. Kopechinskiy, M. Komoń, J. Bissett, G. Szakacs, and C. P. Kubicek. 2005. An oligonucleotide barcode for species identification in *Trichoderma* and *Hypocrea*. *Fungal Genet. Biol.* **42**:813–828.
- Espinel-Ingroff, A., V. Chaturvedi, A. Fothergill, and M. G. Rinaldi. 2002. Optimal testing conditions for determining MICs and minimum fungicidal concentrations of new and established antifungal agents for uncommon molds: NCCLS collaborative study. *J. Clin. Microbiol.* **40**:3776–3781.
- Froslev, T. G., P. B. Matheny, and D. S. Hibbett. 2005. Lower level relationships in the mushroom genus *Cortinarius* (Basidiomycota, Agaricales): a comparison of RPB1, RPB2, and ITS phylogenies. *Mol. Phylogenet. Evol.* **37**:602–618.
- Geiser, D. M., M. del M. Jiménez-Gasco, S. Kang, I. Makalowska, N. Veer-araghavan, T. J. Ward, N. Zhang, G. A. Kuldau, and K. O'Donnell. 2004. FUSARIUM-ID v. 1.0: a DNA sequence database for identifying *Fusarium*. *Eur. J. Plant Pathol.* **110**:473–479.
- Geiser, D. M., M. L. L. Ivey, G. Hakiza, J. H. Juba, and S. A. Miller. 2005. *Gibberella xylarioides* (anamorph: *Fusarium xylarioides*), a causative agent of coffee wilt disease in Africa, is a previously unrecognized member of the *G. fujikuroi* species complex. *Mycologia* **97**:191–201.
- Geiser, D. M., J. I. Pitt, and J. W. Taylor. 1998. Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*. *Proc. Natl. Acad. Sci. U. S. A.* **95**:388–393.
- Gerlach, W., and H. Nirenberg. 1982. The genus *Fusarium*—a pictorial atlas. Mitt. Bundesanst. Land-Forstwirtschaft. Berlin-Dahlem **209**:1–406.
- Gilmore, S. R., T. Gräfenhan, G. Louis-Seize, and K. A. Seifert. 2009. Multiple copies of cytochrome oxidase I in species of the fungal genus *Fusarium*. *Mol. Ecol. Res.* **9**(Suppl. 1):90–98.
- Guarro, J., M. Nucci, T. Akiti, J. Gené, M. D. G. C. Barreiro, and R. T. Gonçalves. 2000. Fungemia due to *Fusarium sacchari* in an immunosuppressed patient. *J. Clin. Microbiol.* **38**:419–421.
- Guarro, J., C. Rubio, J. Gené, J. Cano, J. Gil, R. Benito, M. J. Morandier, and E. Miguez. 2003. Case of keratitis caused by an uncommon *Fusarium* species. *J. Clin. Microbiol.* **41**:5823–5826.
- Gueidan, C., C. R. Villaseñor, G. S. de Hoog, A. A. Gorbushina, W. A. Untereiner, and F. Lutzoni. 2008. A rock-inhabiting ancestor for mutualistic and pathogen-rich fungal lineages. *Stud. Mycol.* **61**:111–119.
- Hall, L., S. Wohlfiel, and G. D. Roberts. 2004. Experience with the MicroSec D2 large-subunit ribosomal DNA sequencing kit for identification of filamentous fungi encountered in the clinical laboratory. *J. Clin. Microbiol.* **42**:622–626.
- Hennequin, C., E. Abachin, F. Symoens, V. Lavarde, G. Reboux, N. Nolard, and P. Berche. 1999. Identification of *Fusarium* species involved in human infections by 28S rRNA gene sequencing. *J. Clin. Microbiol.* **37**:3586–3589.
- Hofstetter, V., J. Miadlikowska, F. Kauff, and F. Lutzoni. 2007. Phylogenetic comparison of protein-coding versus ribosomal RNA-coding sequence data: a case study of the Lecanoromycetes (Ascomycota). *Mol. Phylogenet. Evol.* **44**:412–426.
- Hsiao, C. R., L. Huang, J.-P. Bouchara, R. Barton, H. C. Li, and T. C. Chang. 2005. Identification of medically important molds by an oligonucleotide array. *J. Clin. Microbiol.* **43**:3760–3768.
- Jacobs, A., P. S. van Wyk, W. F. O. Marasas, B. D. Wingfield, M. J. Wingfield, and T. A. Coutinho. 2010. *Fusarium ananatum* sp. nov. in the *Gibberella*

- fujikuroi* species complex from pineapples with fruit rot in South Africa. *Fungal Biol.* **114**:515–527.
32. Jaklitsch, W. M. 2009. European species of *Hypocrea* part I. The green-spored species. *Stud. Mycol.* **63**:1–91.
 33. Kasuga, T., T. J. White, G. Koenig, J. McEwen, A. Restrepo, E. Castañeda, C. da Silva Lacaz, E. M. Heins-Vaccari, R. S. de Freitas, R. M. Zancopé-Oliveira, Z. Qin, R. Negroni, D. A. Carter, Y. Mikami, M. Tamura, M. L. Taylor, G. F. Miller, N. Poonwan, and J. W. Taylor. 2003. Phylogeography of the fungal pathogen *Histoplasma capsulatum*. *Mol. Ecol.* **12**:3383–3401.
 34. Kiehn, T. E., P. E. Nelson, E. M. Bernard, F. F. Edwards, B. Koziner, and D. Armstrong. 1985. Catheter-associated fungemia caused by *Fusarium chlamydosporum* in a patient with lymphocytic lymphoma. *J. Clin. Microbiol.* **21**:501–504.
 35. Koufopanou, V., A. Burt, and J. W. Taylor. 1997. Concordance of gene genealogies reveals reproductive isolation in the pathogenic fungus *Coccidioides immitis*. *Proc. Natl. Acad. Sci. U. S. A.* **94**:5478–5482.
 36. Krulder, J. W. M., R. W. Brimicombe, P. W. Wijermans, and W. Gams. 1996. Systemic *Fusarium nygamai* infection in a patient with lymphoblastic non-Hodgkin's lymphoma. *Mycoses* **39**:121–123.
 37. Landlinger, C., S. Preuner, B. Willinger, B. Haberpursch, Z. Racil, J. Mayer, and T. Lion. 2009. Species-specific identification of a wide range of clinically relevant fungal pathogens by use of Luminex xMAP technology. *J. Clin. Microbiol.* **47**:1063–1073.
 38. Leslie, J. F., and B. A. Summerell. 2006. The *Fusarium* laboratory manual. Blackwell Publishing, Ames, IA.
 39. Liu, Y. J., S. Whelen, and B. D. Hall. 1999. Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Mol. Biol. Evol.* **16**:1799–1808.
 40. Marasas, W. F. O., P. E. Nelson, and T. A. Toussoun. 1984. Toxigenic *Fusarium* species: identity and mycotoxicology. Pennsylvania State University Press, University Park, PA.
 41. Matheny, P. B., Y. J. Liu, J. F. Ammirati, and B. D. Hall. 2002. Using RPB1 sequences to improve phylogenetic inference among mushrooms (*Inocybe*, Agaricales). *Am. J. Bot.* **89**:688–698.
 42. Matheny, P. B. 2005. Improving phylogenetic inference of mushrooms with RPB1 and RPB2 nucleotide sequences (*Inocybe*; Agaricales). *Mol. Phylogenet. Evol.* **35**:1–20.
 43. Mehl, H. L., and L. Epstein. 2007. *Fusarium solani* species complex isolates conspecific with *Fusarium solani* f. sp. *cucurbitae* race 2 from naturally infected human and plant tissue and environmental sources are equally virulent on plants, grow at 37°C and are interfertile. *Environ. Microbiol.* **9**:2189–2199.
 44. Melcher, G. P., D. A. McGough, A. W. Fothergill, C. Norris, and M. G. Rinaldi. 1993. Disseminated hyalohyphomycosis caused by a novel human pathogen, *Fusarium napiforme*. *J. Clin. Microbiol.* **31**:1461–1467.
 45. Migheli, Q., V. Balmas, H. Harak, S. Sanna, B. Scherm, T. Aoki, and K. O'Donnell. 2010. Molecular phylogenetic diversity of dermatologic and other human pathogenic fusarial isolates from hospitals in northern and central Italy. *J. Clin. Microbiol.* **48**:1076–1084.
 46. Naiker, S., and B. Odhav. 2004. Mycotic keratitis: profile of *Fusarium* species and their mycotoxins. *Mycoses* **47**:50–56.
 47. Nelson, P. E., T. A. Toussoun, and W. F. O. Marasas. 1983. *Fusarium* species: an illustrated manual for identification. Pennsylvania State University Press, University Park, PA.
 48. Nucci, M., and E. Anaissie. 2007. *Fusarium* infections in immunocompromised patients. *Clin. Microbiol. Rev.* **20**:695–704.
 49. O'Donnell, K. 2000. Molecular phylogeny of the *Nectria haematococca-Fusarium solani* species complex. *Mycologia* **92**:919–938.
 50. O'Donnell, K., E. Cigelnik, and H. Nirenberg. 1998. Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* **90**:465–493.
 51. O'Donnell, K., C. Gueidan, S. Sink, P. R. Johnston, P. W. Crous, A. Glenn, R. Riley, N. C. Zitomer, P. Colyer, C. Waalwijk, T. van der Lee, A. Moretti, S. Kang, H.-S. Kim, D. M. Geiser, J. H. Juba, R. P. Baayen, M. G. Cromey, S. Bithell, D. A. Sutton, K. Skovgaard, R. Ploetz, H. C. Kistler, M. Elliott, M. Davis, and B. A. J. Sarver. 2009. A two-locus DNA sequence database for typing plant and human pathogens within the *Fusarium oxysporum* species complex. *Fungal Genet. Biol.* **46**:936–948.
 52. O'Donnell, K., H. C. Kistler, E. Cigelnik, and R. C. Ploetz. 1998. Multiple evolutionary origins of the fungus causing Panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proc. Natl. Acad. Sci. U. S. A.* **95**:2044–2049.
 53. O'Donnell, K., H. C. Kistler, B. K. Tacke, and H. H. Casper. 2000. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proc. Natl. Acad. Sci. U. S. A.* **97**:7905–7910.
 54. O'Donnell, K., B. A. J. Sarver, M. Brandt, D. C. Chang, J. Noble-Wang, B. J. Park, D. A. Sutton, L. Benjamin, M. Lindsley, A. Padhye, D. M. Geiser, and T. J. Ward. 2007. Phylogenetic diversity and microsphere array-based genotyping of human pathogenic fusaria, including isolates from the multistate contact lens-associated U.S. keratitis outbreaks of 2005 and 2006. *J. Clin. Microbiol.* **45**:2235–2248.
 55. O'Donnell, K., S. Sink, M. M. Scandiani, A. Luque, A. Colletto, M. Biasoli, M., L. Lenzi, G. Salas, V. González, L. D. Ploper, N. Formento, R. N. Pioli, T. Aoki, X. B. Yang, and B. A. J. Sarver. 2010. Soybean sudden death syndrome species diversity within North and South America revealed by multilocus genotyping. *Phytopathology* **100**:58–71.
 56. O'Donnell, K., D. A. Sutton, A. Fothergill, D. McCarthy, M. G. Rinaldi, M. E. Brandt, N. Zhang, and D. M. Geiser. 2008. Molecular phylogenetic diversity, multilocus haplotype nomenclature, and in vitro antifungal resistance within the *Fusarium solani* species complex. *J. Clin. Microbiol.* **46**:2477–2490.
 57. O'Donnell, K., D. A. Sutton, M. G. Rinaldi, C. Gueidan, P. W. Crous, and D. M. Geiser. 2009. Novel multilocus sequence typing scheme reveals high genetic diversity of human pathogenic members of the *Fusarium incarnatum-F. equiseti* and *F. chlamydosporum* species complexes within the United States. *J. Clin. Microbiol.* **47**:3851–3861.
 58. O'Donnell, K., D. A. Sutton, M. G. Rinaldi, K. C. Magnon, P. A. Cox, S. G. Revankar, S. Sanche, D. M. Geiser, J. H. Juba, J.-A. H. van Burik, A. Padhye, E. J. Anaissie, A. Francesconi, T. J. Walsh, and J. S. Robinson. 2004. Genetic diversity of human pathogenic members of the *Fusarium oxysporum* complex inferred from multilocus DNA sequence data and amplified fragment length polymorphism analyses: evidence for the recent dispersion of a geographically widespread clonal lineage and nosocomial origin. *J. Clin. Microbiol.* **42**:5109–5120.
 59. O'Donnell, K., T. J. Ward, D. M. Geiser, H. C. Kistler, and T. Aoki. 2004. Genealogical concordance between the mating type locus and seven other nuclear genes supports formal recognition of nine phylogenetically distinct species within the *Fusarium graminearum* species complex. *Fungal Genet. Biol.* **41**:600–623.
 60. Özyurt, M., N. Ardiç, K. Turan, Ç. S. Yildiz, O. Özyaral, U. Demirpek, T. Haznedaroğlu, and T. Yurdun. 2008. The isolation of *Fusarium sporotrichioides* from a diabetic foot wound sample and identification. *Marmara Med. J.* **21**:68–72.
 61. Palmore, T. N., Y. R. Shea, R. W. Childs, R. M. Sherry, and T. J. Walsh. 2010. *Fusarium proliferatum* soft tissue infection at the site of plant trauma: recovery, isolation, and direct molecular identification. *J. Clin. Microbiol.* **48**:338–341.
 62. Park, J., B. Park, N. Veeraghavan, K. Jung, Y.-H. Lee, J. E. Blair, D. M. Geiser, S. Isard, M. A. Mansfield, E. Nikolaeva, S.-Y. Park, J. Russo, S. H. Kim, M. Greene, K. L. Ivors, Y. Balci, M. Peiman, D. C. Erwin, M. D. Coffey, A. Rossman, D. Farr, E. Cline, N. J. Grünwald, D. G. Luster, J. Schrandt, F. Martin, O. K. Ribeiro, I. Makalowska, and S. Kang. 2008. *Phytophthora* Database: a forensic database supporting the identification and monitoring of *Phytophthora*. *Plant Dis.* **92**:966–972.
 63. Pennisi, E. 2008. Proposal to "wikify" GenBank meets stiff resistance. *Science* **319**:1598–1599.
 64. Posada, D. 2006. ModelTest Server: a web-based tool for the statistical selection of models of nucleotide substitution online. *Nucleic Acids Res.* **34**:W700–W703.
 65. Pringle, A., D. M. Baker, J. L. Platt, J. P. Wares, J. P. Latgé, and J. W. Taylor. 2005. Cryptic speciation in the cosmopolitan and clonal human pathogenic fungus *Aspergillus fumigatus*. *Evolution* **59**:1886–1899.
 66. Pujol, I., J. Guarro, J. Gené, and J. Sala. 1997. In-vitro antifungal susceptibility of clinical and environmental *Fusarium* spp. strains. *J. Antimicrob. Chemother.* **39**:163–167.
 67. Rakebmann, J. L., U. Bui, K. LaFe, Y.-C. Chen, R. J. Honeycutt, and B. T. Cookson. 2005. Multilocus DNA sequence comparisons rapidly identify pathogenic molds. *J. Clin. Microbiol.* **43**:3324–3333.
 68. Reeb, V., F. Lutzoni, and C. Roux. 2004. Contribution of *RPB2* to multilocus phylogenetic studies of the eukaryotes (Peizozomycotina, Fungi) with special emphasis on the lichen-forming *Acarosporaceae* and evolution of polyspory. *Mol. Phylogenet. Evol.* **32**:1036–1060.
 69. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 70. Schroers, H.-J., K. O'Donnell, S. C. Lamprecht, P. L. Kammeyer, S. Johnson, D. A. Sutton, M. G. Rinaldi, D. M. Geiser, and R. C. Summerbell. 2009. Taxonomy and phylogeny of the *Fusarium dimerum* species group. *Mycologia* **101**:44–70.
 71. Segal, B. H., T. J. Walsh, J. M. Liu, J. D. Wilson, and K. J. Kwon-Chung. 1998. Invasive infection with *Fusarium chlamydosporum* in a patient with aplastic anemia. *J. Clin. Microbiol.* **36**:1772–1776.
 72. Seifert, K. A. 2009. Progress towards DNA barcoding of fungi. *Mol. Ecol. Res.* **9**(Suppl. 1):83–89.
 73. Spiess, B., W. Seifarth, M. Hummel, O. Frank, A. Fabarius, C. Zheng, H. Mörz, R. Hehlmann, and D. Buchheidt. 2007. DNA microarray-based detection and identification of fungal pathogens in clinical samples from neutropenic patients. *J. Clin. Microbiol.* **45**:3743–3753.
 74. Subrahmanyam, A. 1983. *Fusarium laceratum*. *Mykosen* **26**:478–480.
 75. Summerbell, R. C. 2003. *Aspergillus, Fusarium, Sporothrix, Piedraia*, and their relatives, p. 237–498. In D. H. Howard (ed.), *Pathogenic fungi in humans and animals*. Marcel Dekker, Inc., New York, NY.
 76. Summerbell, R. C., and H.-J. Schroers. 2002. Analysis of phylogenetic rela-

- tionship of *Cylindrocarpon lichenicola* and *Acremonium falciforme* to the *Fusarium solani* species complex and a review of similarities in the spectrum of opportunistic infections caused by these fungi. *J. Clin. Microbiol.* **40**:2866–2875.
77. **Sutton, D. A., and M. E. Brandt.** *Fusarium* and other opportunistic hyaline fungi. In J. Versalovic, K. C. Carroll, G. Funke, J. H. Jorgensen, M. L. Landry, and D. W. Warnock (ed.), *Manual of clinical microbiology*, 10th ed., in press. ASM Press, Washington, DC.
78. **Swofford, D. L.** 2002. PAUP*. Phylogenetic analysis using parsimony (and other methods), version 4. Sinauer Associates, Sunderland, MA.
79. **Taj-Aldeen, S. J., J. Gene, I. A. Bozom, W. Buzina, J. F. Cano, and J. Guarro.** 2006. Gangrenous necrosis of the diabetic foot caused by *Fusarium acutatum*. *Med. Mycol.* **44**:547–552.
80. **Taylor, J. W., D. J. Jacobson, S. Kroken, T. Kasuga, D. M. Geiser, D. S. Hibbett, and M. C. Fisher.** 2000. Phylogenetic species recognition and species concepts in fungi. *Fungal Genet. Biol.* **31**:21–32.
81. **Taylor, J. W., E. Turner, J. P. Townsend, J. R. Dettman, and D. Jacobson.** 2006. Eukaryotic microbes, species recognition and the geographic limits of species: examples from the kingdom Fungi. *Philos. Trans. R. Soc. Lond. B* **361**:1947–1963.
82. **Tortorano, A. M., A. Prigitano, G. Dho, M. C. Esposto, C. Gianni, A. Grancini, C. Ossi, and M. A. Viviani.** 2008. Species distribution and *in vitro* antifungal susceptibility patterns of 75 clinical isolates of *Fusarium* spp. from northern Italy. *Antimicrob. Agents Chemother.* **52**:2683–2685.
83. **Walsh, T. J., A. Groll, J. Hiemenz, R. Fleming, E. Roilides, and E. Anaissie.** 2004. Infections due to emerging and uncommon medically important fungal pathogens. *Clin. Microbiol. Infect.* **10**(Suppl. 1):48–66.
84. **Ward, T. J., R. M. Clear, A. P. Rooney, K. O'Donnell, D. Gaba, S. Patrick, D. E. Starkey, J. Gilbert, D. M. Geiser, and T. W. Nowicki.** 2008. An adaptive evolutionary shift in *Fusarium* head blight pathogen populations is driving the rapid spread of more toxicogenic *Fusarium graminearum* in North America. *Fungal Genet. Biol.* **45**:473–484.
85. **Zhang, N., K. O'Donnell, D. A. Sutton, F. A. Nalim, R. C. Summerbell, A. A. Padhye, and D. M. Geiser.** 2006. Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. *J. Clin. Microbiol.* **44**:2186–2190.
86. **Zwickl, D. J.** 2006. Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. Ph.D. dissertation. The University of Texas at Austin, Austin, TX.